

Hypothalamic Hormones

- Hypothalamus → ADH & Oxytocin → neuro-secretory axons → Posterior pituitary
- Hypothalamus → Hormones → network of capillaries (portal system) → Anterior pituitary
- ACTH, TSH, LH, FSH (stimulatory control)
- GH, PRL, MSH (stimulatory and inhibitory control)

■ **General characteristics of hypothalamic hormones:**

TRH, CRH, GHRH, GHIH, GnRH, Dopamine
(DA)

- Small peptides and polypeptides (exception DA) of low M.W
- Needed in very low concentrations (*pg*)
- Have short $t_{1/2}$
- Act on receptors on plasma membrane

■ **TRH=Thyroid Releasing Hormone (Protirelin)**

Tri-peptide, synthetic analogs are available

Effective orally and I.V

Stimulates TSH synthesis and release

MOA: Activation of phospholipase C to increase intracellular IP3 & DAG

Also, TRH has been found to increase PRL release through 2nd messenger Ca⁺⁺

Mainly used:

- As a diagnostic tool (TRH test)
- To treat certain cases of hypothyroidism

Dose: 50 µg I.V, 5 mg orally, maximum response in 15-30 min,
DOA 2-4 hrs

■ CRH=Coricotropin Releasing Hormone

41 a.a peptide stimulates synthesis and release of ACTH, stress \uparrow CRH release

Diagnostic use (CRH test)

■ GHRH (Hexarelin, Sermorelin)

40 a.a peptide, synthetic preparations are available

Diagnostic use and in the management of certain cases of dwarfism (it is given SC)

■ GHIH (Somatostatin)

14 a.a peptide

↓ secretion of GH, ACTH, TSH, Insulin,
Glucagon, Gastrin, Serotonin

Its effects on blood glucose levels are dose
dependent

Low doses → hypoglycemia (↓ glucagon
secretion)

High dose → hyperglycemia (↓ insulin
secretion)

Octreotide (given S.C) & Lanreotide (given I.M)


Synthetic analogs to somatostatin with longer $t_{1/2}$ are mainly used in the management of:

- Acromegaly
- Carcinoid syndrome
- Insulinomas, gastrinomas
- Esophageal varices
- ?? Diabetes mellitus

Major side effects: Gall bladder stone formation and platelet abnormalities

Pituitary Hormones

■ Anterior Pituitary Hormones



■ Posterior Pituitary Hormones

Simple peptides (9 a.a)

ADH (Vasopressin)

Oxytocin

- Hypothalamic hormones regulating the anterior pituitary hormones reach the anterior pituitary through a network of capillaries (portal system) whereas ADH and oxytocin reach the posterior pituitary via neurosecretory axons

Anterior Pituitary Hormones

Hypothalamic lesion or removal → ↓ Ant. Pit H's
except PRL

Hypothalamic stimulation → ↑ Ant. Pit H's except
PRL

■ TSH

↑ T₃ & T₄ through ↑ cAMP, ↑ Iodine uptake

↑ iodination and hydrolysis of thyroglobulin

** diagnostic use

■ ACTH

Derived from larger precursor (Pro-opiocortin)

↑ cortisol release

Undergoes circadian rhythm

Acthar and Cosyntropin (tetracosactrin; Cortrosyn)
are synthetic analogs

Uses:

- Diagnostic use (given I.V or I.M)
- Certain cases of adrenal insufficiency

■ Growth hormone (Somatotropin)

- Species specific
- MOA unclear, its effects believed to be mediated through IGFs (Somatomedins) which are formed in the liver, kidneys, muscles and other tissues
- GH stimulates growth of soft tissues and bones
- ↑ lipolysis
- ↑ gluconeogenesis & ↓ glucose utilization (diabetogenic effect)
- PRL-like activity

■ **Factors ↑ GH release:**

- Sleep, Arginin, Insulin, Hypoglycemia
- β -adrenergic antagonists, Clonidine, Bromocriptine and Levodopa in normal individuals

■ **Factors ↓ GH release:**

- Bromocriptine in acromegalics
- Somatostatin synthetic analogs

■ Disorders affecting GH secreting cells:

Hypersecretion → Gigantism, Acromegaly

R_x

- Surgery
- Somatostatin synthetic analogs
- DA agonists (Bromocriptine; Cabergoline) and
- Pegvisomant (GH-receptor antagonist, given SC, major side effects include abnormal liver enzymes and some reports indicated increased growth of GH-secreting pituitary tumors)

Hyposecretion of GH

- In children it leads to dwarfism manifested by a very short trunk, short neck, shortened arms and legs, average-size hands and feet, broad rounded chest...
- In adults Not so common but it may lead to a higher level of body fat, especially around the waist, anxiety and depression, decreased sexual function and interest, fatigue, less muscle...

Rx of dwarfism → GH replacement therapy

Rx of GH deficiency in adults → loss of weight, good sleep, high protein low carbohydrate diet, exercises ± GH replacement therapy

GH replacement therapy

- GH-replacement therapy with S.C or I.M recombinant human GH preparations:

Somatropin (Humatrope)

Somatrem (Protropin)

- **Mecasermin** (recombinant human IGF-1);
mecasermin rinfabate (recombinant human IGF-1 +IGF binding protein-3=IGFBP-3), given SC in dwarf with IGF-1 deficiency not responding to GH, hypoglycemia is a major side effect

Mecasermin rinfabate is similar to Mecasermin in that both drugs contain recombinant DNA origin insulin-like growth factor 1 (IGF-1). Mecasermin rinfabate however, is already bound to recombinant DNA origin insulin-like growth factor binding protein 3 (IGFBP-3). The binding of IGF-1 to IGFBP-3 is said to extend the half life and reduce the clearance of IGF-1 in patients with growth hormone resistant syndromes and low levels of IGFBP-3 though this may represent <500 patients worldwide

Both mecasermin and mecasermin rinfabate are usually given SC

- Side effects of synthetic rHGH products:

Water retention, the development of antibodies to HGH, insulin resistance and diabetes, hypertension, carpal tunnel syndrome, abnormal bone growth, reduced life span, disturbed insulin metabolism, leukemia, overgrowth of connective tissue, and tumors, ↑ intracranial pressure with papilledema

■ Prolactin (PRL)

Ant. Pit; Placenta

** Dopamine (DA)

** Has GH-like activity

In ♂s PRL increases testosterone production by testes and hence spermatogenesis but \uparrow PRL \rightarrow \downarrow LH & FSH \rightarrow ♂ impotency & infertility

In ♀s:

- Breast development (puberty; pregnancy)

- Lactation

- \uparrow PRL \rightarrow \downarrow LH & FSH (galactorrhea amenorrhea syndrome)

■ Factors/drugs ↑ PRL:

- Pregnancy, sleep, nursing, stress (surgery, exercise)
- TRH, Estradiol, DA antagonists (antipsychotics= phenothiazines and haloperidol; metoclopramide..)
- Methyldopa, reserpine, diazepam, opiates, meclizine, imipramine...

■ Factors/drugs ↓ PRL:

DA agonists (Bromocriptine, pergolide, levodopa)
apomorphine, clonidine , MAO inhibitors (pargyline)

■ Clinical uses to dopamine agonists:

Bromocriptine, Cabergolin...

- Hyperprolactinemia in ♂s and ♀s irrespective of its causes (drug of choice)
- Suppression of lactation
- Acromegaly
- Parkinson's disease
- Cushing's syndrome (Dopamine agonists have been found to inhibit ACTH release)
- DM type II

Dopamine agonists are given orally

Side effects:

Rare, pulmonary fibrosis; confusion; hallucinations; MI...