



Metabolism in erythrocytes

Prof. Mamoun Ahram
Hematopoietic-lymphatic system




- This lecture
- Lippincott's Biochemistry, 8th edition
- The Medical Biochemistry Page
(<https://themedicalbiochemistrypage.org/>)

Carbohydrate metabolism in RBC



- Glycolysis
 - 2,3-bisphosphoglycerate (2,3-BPG)
 - NADH
- Pentose phosphate pathway
 - NADPH






2,3-bisphosphoglycerate (2,3-BPG)

Generation of 2,3-BPG





- 2,3-bisphosphoglycerate (2,3-BPG) is derived in small amounts from the glycolytic intermediate 1,3-bisphosphoglycerate.
- It can re-enter the glycolytic pathway.
 - The erythrocyte loses 2 ATPs.



Effect of 2,3-BPG on Hb





- 2,3-BPG occupies the center of deoxygenated Hb stabilizing it in the T structure.
- When 2,3-BPG is not available (not bound), Hb can be easily converted to the R-structure.




2,3-BPG and HbF



- 2,3-BPG interacts with several groups including His143.
- Fetal hemoglobin (HbF) binds 2,3-BPG much less strongly than HbA.
- Why?



His143 is replaced by a serine in the γ chain.






Glycolysis



Main purpose



- Glycolysis provides
 1. NADH for reduction of methemoglobin (hemoglobin with oxidized Fe^{3+} in heme)
 2. ATP for
 - Modifying sugars and proteins
 - Maintaining membrane asymmetry
 - Functioning of membrane ion pumps
 - Regulating cytoskeletal proteins
 - Maintenance of the discocytic shape, which is critical for the optimal viability and functional capacity.



Phar 6734 – R. L. Johns




Discocyte

Pyruvate kinase isozymes and regulation




- There are two isoenzyme genes of PK and each produces two isoforms:
 - **PKLR gene** produces PKL (liver) and PKR (erythrocytes) using different transcription start sites.
 - **PKM gene** produces PKM1 (muscle and brain) and PKM2 (fetal and most tissues) by alternative splicing.
- Fetal PK isozyme (**PKM2**) has much greater activity than the adult isozymes.
 - Fetal erythrocytes have lower concentrations of glycolytic intermediates including 1,3-BPG and, hence, 2,3-BPG).
 - *Remember: lower 2,3BPG means more Hb in R-state.*

PKLR gene *Different transcription start sites*




PKM gene



Regulation of PKL




- The liver enzyme (PKL) is allosterically regulated:
 - inhibited by ATP, acetyl-CoA, alanine, and long-chain fatty acids and by phosphorylation by protein kinase A.
 - activated by F1,6-BP.
- The liver (PKL) gene is also controlled at the level of synthesis.
 - Increased carbohydrate ingestion induces the synthesis of PKL.



Genetic diseases of PK deficiency

Alterations observed in PK

- The adult erythrocyte PK is virtually inactive.
 - Reduced capacity to make ATP → hereditary hemolytic anemia
- The severity of the disease depends on
 - The degree of enzyme deficiency (5-35%)
 - The ability to produce 2,3-BPG.
- The liver is not affected since expression is stimulated.
- Patients are resistant to malaria.






The pentose phosphate pathway


Two phases of pentose phosphate pathway



- The oxidative generation of NADPH
 - NADPH is generated when glucose 6-phosphate is oxidized to ribulose 5-phosphate.




- The non-oxidative interconversion of sugars



The first step



- The oxidative phase of the pentose phosphate pathway starts with the dehydrogenation of glucose 6-phosphate by glucose 6-phosphate dehydrogenase (G6PD).





- G6PD is highly specific for NADP^+ , relative to NAD^+
- The reaction is irreversible and is the rate-limiting reaction.
- High levels of NADP^+ stimulate the reaction .

Oxidative stress and glutathione

- Oxidative stress within cells is controlled by the action of glutathione (GSH).
- GSH reduces peroxides via glutathione peroxidase.
- GSH is regenerated via NADPH-dependent glutathione reductase.
- The PPP in erythrocytes is the only pathway to produce NADPH.


PPP consumes almost 10% of glucose by erythrocytes.



Low GSH levels



- The inability to maintain reduced glutathione in RBCs leads to increased accumulation of peroxides, predominantly H_2O_2 , resulting in weakening of the cell membrane due to:
 - peroxidizing membrane lipids leading to hemolysis
 - oxidizing proteins including hemoglobin (to methemoglobin) and membrane proteins, insolubilizing them, and forming Heinz bodies







Glucose-6-phosphate dehydrogenase deficiency

G6PD deficiency



- Glucose-6-phosphate dehydrogenase (G6PD) deficiency is a group of heterogeneous disease with significantly reduced activity.
 - **Hemolytic anemia**
 - particularly after the administration of drugs, during infections and in the neonatal period (jaundice)
- Deficiency of G6PD is most prevalent in individuals of African, Mediterranean, and Oriental ethnic origins.
- It is the most common enzyme deficiency worldwide.
- G6PD gene is located on the X chromosome.
 - **Inheritance of G6PD deficiency is sex-linked.**






Frequency of G6PD deficient males %



Polymorphic G6PD variants



G6PD mutations



- Several hundred G6PD genetic variants have been identified, but most have no clinical symptoms.
- Almost all G6PD deficiency variants are caused by point mutations in the gene.
 - These mutations mainly alter the kinetic properties, stability, or binding affinity to NADP⁺ or G6P.
- No large deletions or frameshift mutations. Why?

The four classes of G6PD deficiency



- G6PD B (Normal)

- Abnormal G6PDs

- Class I: the most severe and rare.

- Class IV: no clinical symptoms

- G6PD A- (group III or class III)

- Among persons of African descent

- It is caused by a single amino acid substitution that decreases enzyme (protein) stability, but has 5-15% of normal activity.

- The disease is moderate.

- G6PD Mediterranean (group II or class II)

- Severe

- The enzyme has normal stability, but negligible activity.

Class	Clinical symptoms	Residual enzyme activity
I	Very severe (chronic hemolytic anemia)	<2%
II	Severe (episodic hemolytic anemia)	<10%
III	Moderate	10%-60%
IV	None	>60%

Class II vs. class III




G6PD A- (class III):

Moderate, young RBCs contain enzymatic activity. Unstable enzyme, but kinetically normal

G6PD Mediterranean (II)

Enzyme with normal stability but low activity (severe). Affect all RBCs (both young and old)

Although the activity of the normal enzyme declines as red cells age, even the oldest cells have a sufficient level of activity to provide protection against oxidative damage and hemolysis.



By contrast, very few *G6PD Mediterranean* red cells have sufficient enzyme activity to prevent oxidative damage, whereas a substantial fraction of young *G6PD A⁻* red cells are able to provide protection.

Inducers of G6PD deficiency symptoms



- Oxidant drugs
 - Antibiotics, anti-malarial, and anti-pyretics (not acetaminophen)
- Fava beans (favism)
 - Fava beans are presumed to cause oxidative damage.
 - Substances capable of destroying red cell GSH have been isolated from fava beans (fool).
 - Favism is most common in persons with G6PD class II variants, but rarely can occur in patients with the G6PD A- variant.
- Infection
 - The most common inducer due to production of free radicals.

Connection to malaria

- Several G6PD deficiencies are associated with resistance to the malarial parasite, *Plasmodium falciparum*, among individuals of Mediterranean and African descent.
- The basis for this resistance is the weakening of the red cell membrane (the erythrocyte is the host cell for the parasite) such that it cannot sustain the parasitic life cycle long enough for productive growth.

