

Past papers diveded by lectures

Antidepressant drugs

Orthostatic hypotension is a side effect of?

- A. Bupropion
- B. Amitriptyline
- C. Phenezine
- D. Mirtazapine
- E. None of the above

Answer: B

Which of the following drugs is best used in atypical depression

Answer: Selegiline

Mao inhibitor

- wrong matching :

Answer: paroxetine >> decreased body weight

Antiphscosis

Which of the following is a cause of schizophrenia only?

A. Too much serotonin

B. Too much dopamine

C. Too much norepinephrine

D. Decreased dopamine levels

E. All of the above

Answer: B

The drug which is used in refractory psychosis:

Answer: Clozapine (last resort)

أتوقع مش مطلوب

wrong matching :

Answer: Clonazpine:relapse take weeks

Here's the proper matching:

- Clozapine is an atypical antipsychotic used especially in treatment-resistant schizophrenia.
- One of its unique features is that relapse after stopping clozapine can happen very quickly, often within a few days, not weeks.
- This is in contrast to other antipsychotics, where relapse might take longer after discontinuation.

Correct Matching:

- Clozapine: Relapse occurs within days after stopping the drug.

This is one reason why adherence to clozapine is so crucial, and sudden discontinuation is avoided unless absolutely necessary.



Increasing the dose of haloperidol,
decreases the side effect of?

A. Tardive dyskinesia

B. Akathisia

C. Acute dystonia

D. Orthostatic hypotension

Answer: A

أتوقع السؤال خطأ

Which of the following drugs is correctly matched with its side effects

A. Lithium- hyperthyroidism

B. Lithium- diabetes insipidus

C. Olanzapine- agranulocytosis

D. Clozapine- diabetes mellitus

E. Quetiapine- elevation of prolactin

Answer: B

Bipolar drugs

163. Which of the following drug is correctly matched with its side effect?

A) Lithium - diabetes insipidus

B) Olanzapine-Agranulocytosis

ANSWER : A

Lithium toxicity is associated with decreased Na^+ , so what happens in the case of increased Na^+ ? (This question was deleted)

- A. Increased lithium reabsorption
- B. Increased lithium clearance
- C. Decreased lithium reabsorption
- D. Lithium concentration is not affected
- E. None of the above

Answer: C

ANSWER: B

. Which of the following drugs is correctly matched with its side effects

- A. Lithium- hyperthyroidism
- B. Lithium- diabetes insipidus
- C. Olanzapine- agranulocytosis
- D. Clozapine- diabetes mellitus
- E. Quetiapine- elevation of prolactin

Answer: B

37. What of the following is a good choice if a doctor wants to use a general anaesthetic with short and fast induction time:

Answer: propofol

343. The most used general anesthetic in operations with fast and short induction ?

- A) propofol.
- B) Thiopental.
- C) Etomidate.
- D) Ketamine.
- E) Halothane.

ANSWER : A

Local anaesthetic drugs

344. One of the following is categorized as local anesthetic :

- A) halothane.
- B) Nitrous oxide.
- C) Mepivacaine.
- D) Thiopental.
- E) Ketamine.

ANSWER : C

Parkinson and Alzheimer Drugs

348. Patients with Alzheimer disease which of the following drug Is good to treat their condition ?

- A) NMDA receptor antagonist.
- B) Acetylcholine receptor antagonist.
- C) Giving acetylcholine esterase.
- D) MAOI's.
- E) Levodopa

ANSWER : A

349. Patient with Parkinson's disease are advised to take ?

- A) dopamine.
- B) Levodopa.
- C) Acetylcholine.
- D) SSRI.
- E) Anti-acetylcholine esterase.

ANSWER : B

350. In Parkinson disease we give levodopa as a drug instead of dopamine, the reason behind

Metabolism and BBB

Epilepsy drugs

38. A patient is suffering from epilepsy and not responsive to traditional drugs, which of the following drugs can be used in this case:

- A. Cl⁻ channel blocker
- B. GABA reuptake inhibitors
- C. GABA antagonists
- D. Glutamate agonists

Answer: B

351. A 22-year-old female presented to ER suffering of epileptic attack, which one of the following drugs you should not give ?

- A) Sodium channels blocker.
- B) calcium channels blocker.
- C) GABA reuptake enhancer.
- D) Phenytoin.
- E) All of the above can be given to epileptic patient.

ANSWER : C

345. Patients with attention-defect hyperactivity disorder should take which one of the following ?

- A) amphetamines.
- B) GABA reuptake inhibitor.
- C) Serotonin antagonist.
- D) D2 blocker.
- E) SSRI.

ANSWER : A

346. A patient presented to your clinic suffering from narcolepsy, the proper drug to use is ?

- A) MAOI.
- B) Amphetamine.
- C) SNRI.
- D) SSRI.
- E) Serotonin antagonist.

ANSWER : B

34. Patient with depression, for treating narcolepsy:

- A. Dopamine agonist
- B. Serotonin agonist
- C. Serotonin antagonist
- D. Dextroamphetamine

Answer: D

تم بحمد الله ،،،،،

لا تنسونا من صالح الدعاء 

أكثرُوا الدعاء لنصرة المسلمين في غزة