



FLUID MANAGEMENT AND BLOOD TRANSFUSION

Dr. Mahmoud Almustaf, M.D
FACHARZT.



Body Fluid Compartments

- TBW: 55-60% of the BW in men and 45-50% in



Body Fluid Composition in Age Groups

AGE	TBW AS % OF TOTAL BODY WEIGHT
Neonate	80
6 months	70
1 year	60
Young adult	60
Elderly	50

Composition of Body Fluid Compartments

• Ion (mmol/L)	Plasma (mmol/L)	ICF
• Na ⁺	143	9
• K ⁺	5	135
• Ca ²⁺	1.3	<0.8
• Mg ²⁺	0.9	25
• Cl ⁻	103	9
• HCO ₃ ⁻	24	9
• HPO ₄ ²⁻	0.4	74
• Sulphate ⁻	0.4	19
• Proteinate ⁻	1.14	64

ESSENTIAL PRINCIPLES

Osmolarity and Osmolality

- These are ways of quantifying how much of a solute is dissolved in a solution.
- ***Osmola(R)ity*** No. of osmoles of solute particles per unit VOLUME of solution and has units **osmoles/litre**. In the body we use **milliosmole**
- ***Osmola(L)ity*** No. of osmoles of solute particles per unit WEIGHT of solvent and has units **osmoles/kilogram**.

Plasma Osmolality

Plasma osmolality = $2 (\text{Na} + \text{K}) + \text{glucose} + \text{urea}$

= $2 (137 + 4.0) + 5.0 + 4$


= **291** mosmol/kg H₂O

Glu: /18

Urea: /2.8

Tonicity

- A way of describing the relative solute concentrations of **two solutions** which are separated by a selectively-permeable membrane (often called a **semi-permeable membrane**).



WHAT IS THE 'NORMAL'
DAILY INTAKE AND
— OUTPUT OF FLUID AND —
ELECTROLYTES?

OUTPUT, INTAKE

- Input: Oral, Enteral, Intravenous
- Output: ‘Sensible’: that it is easily seen and measured e.g. urine output and losses from the gastrointestinal tract.
- ‘Insensible’: not seen and not easy to quantify e.g. sweat, and water vapor in exhaled gases.

OUTPUT

Urine

1500 ml

Gastrointestinal(faeces)

200 ml

Skin(sweat)

400 ml

Respiratory

400 ml

Total

2500

INTAKE

Drinking

1500 ml

Eating

750 ml

Metabolism

250 ml

Total

2500

- The above volumes do not contain just water but also **electrolytes**....

ELECTROLYTE	DAILY LOSS / REQUIREMENT	75 Kg PERSON PER DAY
Sodium	1-1.5 mmol/kg	75 - 112.5 mmol
Potassium	1-1.5 mmol/kg	75 – 112.5 mmol
Magnesium	0.1-0.2 mmol/kg	7.5 – 15 mmol
Calcium	0.1-0.2 mmol/kg	7.5 – 15 mmol
Chloride	0.07-0.22 mmol/kg	5.25 – 16.5 mmol
Phosphate	20-40 mmol/kg	1500 – 3000 mmol

Maintenance and Deficit

- Maintenance Vs deficit
 - Rule of 4 /2/ 1
 - Ex. 70 kg patient
 - 1st 10 kg: $10 \text{ kg} * 4 \text{ ml} = 40 \text{ ml / kg}$
 - 2nd 10 kg: $10 \text{ kg} * 2 \text{ ml} = 20 \text{ ml / kg}$
 - 3rd 10 kg: $50 \text{ kg} * 1 \text{ ml} = 50 \text{ ml / kg}$
- Total=110ml/kg
- Ex. Fasting for 10 hr without any intake: $10 * 110 = 1100 \text{ ml}$

Allowable Blood Loss (ABL)

- $EBV = \text{weight (kg)} * \text{Average blood volume}$
- $\text{Allowable Blood Loss} = [EBV * (H_i - H_f)] / H_i$
- Where:
 - $EBV = \text{Estimated Blood Volume}$
 - $H_i = \text{initial hemoglobin (Hct)}$
 - $H_f = \text{final hemoglobin (Hct)}$
- Normal Hct Values
 - Men 42-52%
 - Women 37-47%

Estimated Blood Volume (EBV)

- Men 75 ml / kg
- Women 65 ml / kg
- Infants 80 ml / kg
- Neonates 85 ml / kg
- Premature Neonates 96 ml / kg

INTRAVENOUS FLUIDS

Types

- Three main types
 - Crystalloids
 - Colloids
 - Blood products

Crystalloids ...



- Solutions that contain a combination of **water and electrolytes**.
- Divided into "***balanced***" salt solutions (e.g. Ringer's lactate) and ***hypotonic*** solutions (e.g. D5W).

- Classified into three groups based on their predominant use

- Replacement Solutions

- Maintenance Solutions

- Special Solutions

Replacement Solutions

- Used to replace **ECF**
- All **isotonic**, usually **replace losses** that involves both water and electrolytes
- Have a $[\text{Na}^+]$ similar to that of the ECF which effectively **limits** their fluid **distribution** to the **ECF** compartment.
- Distributes between the ISF $\frac{3}{4}$ and the plasma $\frac{1}{4}$ in proportion to their volumes

Ringer's Lactate (Hartman's)

- $\text{Na}^+ = 131$ mmol/L
- $\text{Cl}^- = 111$ mmol/L
- Lactate = 29 mmol/L
- $\text{K}^+ = 5$ mmol/L
- $\text{Ca}^{++} = 2$ mmol/L
- $\text{PH} = 6.5$
- Osmolality = 279 mosm/L
- Potential problem = potassium may accumulate,

Maintenance Solutions

- **Isosmotic** as administered but not necessarily isotonic
- Usually used when the loss involves mainly pure water
- Ex. **D5W, Normal Saline**

Normal saline (0.9% saline solution)

- 9 g of NaCl/L water
- 154 mmol/L sodium
- 154 mmol/L chloride
- Osmolality = 308 mosm/L
- PH = 5.0
- Potential problem = hyperchloraemic metabolic acidosis, more likely with renal insufficiency

Special Solutions

- Hypertonic (3%) saline.... hyponatremia
 - 30 gm NaCl, 1027, 4.5 to 7.0
- Half normal saline.... hypernatremia
 - 77 meq/L
- 8.4% Bicarbonate solution... acidosis
- Mannitol 20%....brain oedema, pulmonary oedema

colloids...



Starches

Colloids

- Colloid: a **large molecule** that **does not diffuse** across semipermeable membranes (capillary)
- Exerts an **osmotic pressure** in the blood, causing fluid to remain within the vascular system. The result is an increase in intravascular volume.

- Two categories of colloid may be defined:
 - **Natural** (e.g. human albumin)
 - **Artificial** (e.g. gelatins, dextran and hydroxyethyl starches [HES]).

Albumin

- Half-life ($t_{1/2}$) = 1.6 hours in plasma
- Stays within the intravascular space unless the capillary permeability is abnormal
- 5% solution - isotonic; 20% solutions - hypertonic
- Expands volume 5x its own volume in 30 minutes
- Side effects - volume overload, fever (pyrogens in albumin), defects of haemostasis

Dextran

- High MW polysaccharide
- Dextran 40 - MW 40,000
- Dextran 70 - MW 70,000
- 10% solution in NS or D5W
- Side effects: anaphylaxis, coagulopathy, renal failure
- Dose: limit to 20 ml/kg/day
- Used as antiaggregant in patients undergoing vascular and microvascular surgical procedures

PERIOPERATIVE BLOOD TRANSFUSION

Purpose of Infusion of Fluids and Blood Products

- Maintain organ transfusion
- Normal blood pressure and heart rate, normal mental status (in non-comatose patients), normal oxygen saturation, normal urine output, well perfused extremities

Blood Products

- Whole blood
- Packed Red Blood Cells
- Platelets
- Fresh Frozen Plasma
- Cryoprecipitate
- Human Albumin

BLOOD BANK PRACTICES

Preparation of Blood Components


- Blood donors:
 - Approximately 17 million units of blood are donated in Europe each year.
 - Each donor is interviewed for medical history of known infectious diseases
 - Each unit is screened for antibodies to:
 - Syphilis
 - Hepatitis B and C
 - HIV 1 and 2
 - +/- CMV

Centrifugation


- Collect 500 mL whole blood
- Divert the first 40 mL to reduce risk of bacterial contamination from donor skin
- The 40 mL are used for donor unit testing
- Blood is centrifuged and separated into 3 parts:
 - ◆ Red Blood Cells
 - ◆ Plasma
 - ◆ Buffy coat

Process for Preparing Blood Components from Donated Units


Step 1 - Whole Blood Separation



Step 2 - RBC



- The Buffy coat units from four donors are further processed to separate the platelets
- The red blood cell and platelet components are



- Separated of blood components by 1 unit of **Whole blood**:
- **PRBCS**(hematocrit 70%):
 - 250 mL+saline preservative=350 mL.
 - 1–6°C.
 - May be frozen in a hypertonic glycerol solution for up to 10 years(rare phenotypes)

- A preservative–anticoagulant solution is added. The most commonly used solution is **CPDA-1**:
- **C**itrate as an anticoagulant (by binding calcium)
- **P**hosphate as a buffer
- **D**extrose as a RC energy source
- **A**denosine as a precursor for ATP synthesis.
- 35 days
- AS-1 (Adsol) or AS-3 (Nutrice) extends the shelf-life to 6 weeks.
- ADSOL (Adenine, glucose, mannitol and sodium chloride)
- NUTRICE (Adenine, glucose, citrate, phosphate and NaCl)

- **Platelet:**

- 50–70 mL.
- 20–24°C for 5 days.

- **Plasma:**

- The remaining plasma supernatant is further processed and frozen to yield fresh frozen plasma; rapid freezing helps prevent inactivation of labile coagulation factors (V and VIII). Slow thawing of fresh frozen plasma yields a gelatinous precipitate (cryoprecipitate) that contains high concentrations of Factor VIII and fibrinogen.
- 200 mL.
- Once thawed it must be transfused within 24 h.

**BUT BEFORE EVERYTHING
THE BLOOD BANK HAS
TESTS TO COMPARE THE
BLOOD OF THE DONOR TO
THE BLOOD OF THE
RECIPIENT**

So you must know the blood groups

Blood Groups

- At least 20 separate blood group antigen systems are known; fortunately, only the ABO and the Rh systems are important in the majority of blood transfusions.

The ABO System

- Simply speaking, the chromosomal locus for this system produces two alleles: A and B. Each represents an enzyme that modifies a cell

Incidence	Naturally Occurring Antibodies in Serum	Type
45%	Anti-B	A
8%	Anti-A	B
4%	—	AB (Universal recipient)
43%	Anti-A, anti-B	O (Universal donor)

COMPATIBILITY TESTING

Intra-operative Transfusion Practices

Packed Red Blood Cells

- Ideal for patients requiring RCs but not volume replacement (eg, anemia pt in compensated CHF).
- Hgb 7-8 g/dL (<6, most people require blood >10 most people do not)
- Each unit raise Hgb by 1g/dl
- 170- μ m filter to trap any clots or debris.
- Warming to 37°C during infusion.
- Hypothermia and low levels of 2,3-diphosphoglycerate (2,3-DPG) in stored blood can cause a marked leftward shift of the hemoglobin–oxygen dissociation curve
- ABO-compatible units are mandatory.



Fresh Frozen Plasma

- FFP contains all plasma proteins, including all clotting factors.

- Indications:

- Isolated factor deficiencies.
- Reversal of warfarin therapy.
- Coagulopathy associated with liver disease
- CABG, bleeding+NL ACT.
- Massive blood transfusions.
- Antithrombin III def.



- The initial therapeutic dose is usually 10–15 mL/kg
- ABO-compatible units are mandatory.
- Coagulation factors INR 1.4-1.6 (INR>1.6, most people require FFP transfusion for major surgery; INR<1.4, most people do not require)

Platelets

- Thrombocytopenia or dysfunctional platelets .
- Surgery or invasive procedures: $70,000 \times 10^9$
- Vaginal delivery and minor surgical procedures: $50,000 \times 10^9/L$.
- Each unit expected to increase the count by $10,000\text{--}20,000 \times 10^9/L$.
- ABO-compatible platelet transfusions are desirable but not necessary



Cryoprecipitate


- Each unit (15 ml) contains fibrinogen 150 mg, factor VIII 100 units, von Willebrand factor (vWF) (100 units)
- DIC, hemophilia A, von Willebrand disease, quick reversal of thrombolytic therapy
- Fibrinogen (most people require cryoprecipitate for major surgery if fibrinogen < 1 g/dL)



CELL SAVER



CELL SAVER



LEVEL 1 (A) INFUSION PUMP



Complications of Blood Transfusion

- Hemolytic reactions
 - Acute Vs. Delayed
- Febrile Non hemolytic reactions
- Transfusion Related Acute Lung Injury (TRALI)
- Infectious complications

TRALI

- ARDS following blood transfusion
- High morbidity ... mechanical ventilation
- Lung injury is generally transient with PO₂ levels returning to pretransfusion levels within 48 -96 hours and CXR returning to normal within 96 hours.
- Mortality rate, often approximated at 5 to 10%
- Treatment as ARDS

APPROACHES TO FLUID MANAGEMENT

The “Classic” Approach

- Step 1: Calculate Ongoing Maintenance Requirements

4/2/1 rule: 4 cc/kg/hr for the first 10 kg, 2 cc/kg/hr for the second 10 kg, and 1 cc/kg/hr for every kg above 20.

Ex. 70kg $\rightarrow 10 * 4 \text{ ml} = 40 \text{ ml}$

$10 * 2 \text{ ml} = 20 \text{ ml}$

$50 * 1 \text{ ml} = 50 \text{ ml}$

110ml / **hour** given each hour as
oral or enteral intake is stopped

Ex. 15kg $\rightarrow 10 * 4 \text{ ml} = 40 \text{ ml} + 5 * 2 \text{ ml} = 10 \text{ ml} = 50 \text{ ml} / \text{hour}$

- Step 2: Calculate Preoperative Fluid deficit
Maintenance * the time without intake what so ever

Ex. 70kg fasting for 10 hours pre-operatively

$$M = 110\text{ml/hour} * \text{time } 10 \text{ hour} = 1100 \text{ ml}$$

- Step 3: Calculate Anticipated Surgical Fluid Losses
- Minimal tissue trauma (ex. herniorrhaphy): 0-2 cc/kg/hr
- Moderate tissue trauma (ex. cholecystectomy): 4-6 cc/kg/hr
- Severe tissue trauma (ex. bowel resection): 8-10 cc/kg/hr
- Ex. 70kg undergoing major laparotomy →
 $10 \text{ ml} * 70 \text{ kg} = 700 \text{ ml/hour}$ as long as surgery is going on

- Step 4: Adjust for Blood Losses
- A common recommendation is to give 3-4 cc of crystalloid for every 1 cc of blood loss
- Remember to add up suction volume, lap pads (100-150 cc each if fully soaked) and 4x4 small pads (10 cc each if fully soaked)

Ex. In the 1st hour of laparotomy there was 200ml of pure blood in the suction jar, 2 fully soaked lap pads, and 10 fully soaked small gauzes

$200 \text{ ml} + 2 * 150 \text{ ml} + 10 * 10 \text{ ml} = 600 \text{ ml}$ (in that hour) to be replaced with either $600 * 4$ of cryst. or -if indicated- 600 ml of blood

- 1st hour = Maintenance + $\frac{1}{2}$ Deficit + Blood loss + Ongoing loss
- 2nd hour = Maintenance + $\frac{1}{4}$ Deficit + Blood loss + Ongoing loss
- 3rd hour = Maintenance + $\frac{1}{4}$ Deficit + Blood loss + Ongoing loss
- Maintenance continued post-operatively as long as fasting
- Blood loss replaced as long as there is bleeding
- Ongoing loss as long as the surgery continues

THANK YOU
