




Gynecology & Obstetrics
For undergraduates

هدية
مجانية

Notes on the Board of Dr. Nadine's Lectures

6th Edition
2025



Notes of Dr. Nadine Alaa Sherif
Professor of Obstetrics & Gynecology
Faculty of Medicine – Cairo University

Written by:
Dr. Reem Abdelhakim
Rheumatology resident in Kasr Al Ainy

www.nadine-alaa-sherif.weebly.com



*This work is dedicated to the soul of my father,
my beloved mother, my unique brother, and ...
All the future doctors, whom I truly believe in
their creativity, intentions and potentials .*

Nadine Alaa Sherif



Website



YouTube channel



Telegram channel



Facebook Page

List of Abbreviations

AFE : Amniotic Fluid Embolism	IAP: Intra abdominal pressure	O/E : On Examination
AFI : Amniotic Fluid Index	ICHge : Intracranial hemorrhage	OF: Occipito-frontal
AGA : Average for gestational age	II LNs : Internal iliac lymph nodes	OGTT : Oral Glucose Tolerance test
AIS : Androgen insensitivity syndrome	II vessels : Internal Iliac vessels	OHSS: Ovarian hyperstimulation syndrome
ANC : Antenatal care	IIA : Internal iliac artery	PAPP-A : Pregnancy Associated Plasma Protein A
ART : Assisted Reproductive Technology	IIV : Internal iliac vein	PAS : Placenta Accreta Spectrum
ASCUS : Atypical squamous cells of undetermined significance	IMB : Intermenstrual Bleeding	PCT : Post coital test
AUB : Abnormal Uterine Bleeding	IPC: Intrapartum care	PE : Preeclampsia
BAD : Bis-acromial diameter	IPHge : Intra Peritoneal Hemorrhage	PEB : Premenstrual Endometrial Biopsy
BCT : Benign cystic teratoma	IUFD : Intrauterine fetal death	PG: Primigravida / Pelvic Grip
BL : Broad Ligament	IUI : Intrauterine insemination	POI : Premature ovarian insufficiency
BLM : Broad ligament myoma	IUP : Intraurethral pressure	POP: Pelvic Organ Prolapse
BMD: Bimastoid diameter	IVP: Intravesical pressure	PPC : Post Partum Care
BPD : Biparietal diameter	LEEP : Loop Electrosurgical Excision Procedure	PPT Labor : Precipitate labor
BPB: Biophysical profile	LGA : Large for gestational age	PROM : Premature Rupture of membranes
BTB : Bitrochanteric diameter / Bitemporal diameter	LGV: Lymphogranuloma Venereum	PUL : Pregnancy of Unknown Location
BVs : Blood vessels	LLETZ : Large loop excision of the transformation zone	ROD : Right Oblique Diameter
CIA : Common Iliac Artery	LMA : Left Mento Anterior	ROM : Rupture of membranes
CILNs : Common Iliac Lymph nodes	LMP :Last menstrual period / Low malignant potential	RPL : Recurrent Pregnancy Loss
CPR: Cardiopulmonary resuscitation	LMWH : Low Molecular Weight Heparin	RSA : Right scapula – anterior
DFMC : Daily fetal movement count	LNG-IUD : Levonorgestrel - Intrauterine Device	SGA : Small for gestational age
DL : Diagnostic laparoscopy	LOA : Left Occipito Anterior	SIDS : Sudden infant death syndrome
DMA : Direct Mento Anterior	LPD : Luteal phase defect	SMB: Submento-bregmatic
DO : Detrusor overactivity	LSA : Left scapula –anterior	SMM : Submucous myoma
DOA : Direct Occipito Anterior	LSIL : Low grade squamous intraepithelial lesion	SMV: Submento-vertical
DOP : Direct Occipito Posterior	MA : Mento-anterior / Monoamniotic	SOB: Suboccipito-bregmatic
DZT : Dizygotic Twin	MAC : Methotrexate, Actinomycin – D, Cyclophosphamide	SOF : Suboccipito-frontal
EH : Endometrial Hyperplasia	MC : Monochorionic	SP: Symphysis Pubis
ET : Endometrial Thickness	MCL : Midclavicular line	TD : Tubal Disconnection
EILNs : External Iliac Lymph nodes	MCT : Malignant cystic teratoma	TDF : Testicular differentiation factor
FCA : Fetal congenital anomalies	MDIF : Mullerian duct inhibiting factor	TENS : Transcutaneous Electrical Nerve Stimulation
FFP : Fresh Frozen Plasma	MH : Metropathia haemorrhagica	TOA : Tubo-ovarian abscess
FG : Fundal Grip	MP : Mento-posterior	TOC : Tuboovarian cyst
FGM : Femal genital mutilation	MRKH : Mayer Rokitansky Küster Hauser syndrome	TOT : Transobturator tape
FL: Fundal Level	MSAFP : Maternal serum Alpha Fetoprotein	TTTT : Twin To Twin Transfusion syndrome
FU : Follow Up	MSH : Melanocyte stimulating hormone	TVT : Tension-free vaginal tape
GI : Granuloma Inguinale	MST: Malignant solid teratoma	UAE: Uterine artery embolisation
HIFU: High Intensity Focused Ultrasound	MTX : Methotrexate	UG : Umbilical Grip
HMB : Heavy Menstrual Bleeding	MV: Mento-vertical	VaIN : Vaginal intraepithelial neoplasia
HPL : Human Placental Lactogen	MZT : Monozygotic Twin	VIN : Vulvar intraepithelial neoplasia
HSIL : High grade squamous intraepithelial lesion	NST : Non stress test	VVF : Vesicovaginal fistula
HSM : Hepatosplenomegaly	NT : Nuchal Translucency	
	NTD : Neural Tube Defect	

Index

Gynecology

Physiology of menstrual cycle -----	7
Amenorrhoea -----	8
Puberty & Menopause -----	9
Anovulation ,LPD, PCO , Hirsutism & Hyperprolactinemia -----	10
Infertility & ART -----	11
FSD & FGM -----	12
Fibroid -----	13
Endometriosis & Adenomyosis -----	14
AUB , Dysmenorrhea , PMS -----	15
Functional cysts of the ovary -----	16
Benign conditions of the vulva & vagina -----	16
Contraception -----	17
Pelvic Organ Prolapse -----	18
Urinary Incontinence & Rectovaginal fistula -----	19
Lower genital tract infections -----	20
Upper genital tract infections -----	21
STDs -----	23
Scheme for oncology-----	24
Endometrial Carcinoma -----	25
Cancer Cervix -----	26
Benign Ovarian Tumors -----	27
Low Malignant potential (LMP) ovarian tumors -----	28
Malignant Ovarian Tumors -----	29
Tumors of vulva & vagina -----	30
Anatomy of female genital tract -----	31
Sexual differentiation & abnormalities of female genital system ---	34
Endoscopy in Gynecology -----	35

Obstetrics

Abortion -----	37
Ectopic pregnancy -----	38
GTDs -----	39
APHge / Intrapartum Hge -----	40
Complications of 3rd stage of labor / PAS -----	41
DIC / AF embolism / maternal collapse & mortality -----	42
Anatomy of female pelvis & fetal skull -----	43
Normal Labor -----	44
Occipito posterior / Face & Brow presentations -----	45
Breech presentation -----	46
Shoulder presentation / cord presentation & prolapse / complex presentation -----	47
Multifetal gestation -----	48
Operative vaginal delivery -----	49
CS / Episiotomy / Analgesia & Anesthesia in labor -----	50
Fetal surveillance (Antepartum & Intrapartum) -----	51
Abnormal labor & CPD -----	52
Premature Rupture of Membranes / Amniotic fluid disorders -----	54
Prematurity & Postmaturity -----	55
SGA & LGA & IUFD -----	56
Fetal & neonatal asphyxia -----	57
Fetal birth injuries -----	58
Puerperium & Puerperal pyrexia -----	59
Puerperal sepsis -----	60
RH isoimmunization -----	61
Hypertension with pregnancy -----	62
DM with pregnancy -----	63
UTI , Cardiac diseases & Seizures with pregnancy -----	64
Anemia , Thyroid disorders ,VTE & PE with pregnancy -----	65
GIT disorders & hepatic disorders with pregnancy -----	66
Prenatal diagnosis of congenital anomalies -----	67
Fertilization, implantation -----	68
Fetal circulation -----	69
Placenta , cord -----	70
Diagnosis of pregnancy / Physiological changes during pregnancy -----	71
ANC -----	71
High risk pregnancy -----	72

Instruments in OBGYN 73-74

Clinical History taking 76-81

GYNECOLOGY

Physiology of menstrual cycle

Hypothalamus :
GnRH




Pituitary :
FSH / LH

Ovary:
E₂ / P

Endometrium :
Glands

Cervix:
Mucus

Vagina:
Cells



NB:

- * 2 Cell theory: theca & granulosa cells for E₂ synthesis (steroidogenesis).
- * Other catalysts as Inhibin - - FSH, Activin ++FSH
- * PGs of secretory endometrium & dysmenorrhea
- * PMS & ovulation (PRG effect)

Amenorrhea

Def:


Et:

- Outflow (cryptomenorrhea)
- Uterus
- Ovaries
- Ant pituitary
- Hypothalamus

Cl pict for each

Investigations

Ttt for each



Approach to a case of amenorrhea :

1) History (Et) :

drugs , 1^{ry} , 2^{ry} , PPH , D&C , ...

2) Examination (cl.picture):

2^{ry} sexual ch / local...

3) Inv (diagnosis) :

For 1^{ry} amenorrhea

In presence of 2^{ry} sex. ch.

↓
US

In absence of 2^{ry} sex. ch.

↓
Hormonal profile
FSH/LH

Others

(karyotyping / laparoscopy / hysteroscopy /....)

4) ttt (medical or surgical) :

depending on etiology

Puberty (11-12yrs)


(Non reproductive → reproductive)

Due to maturation of pulsatility of hypothalamus

Normal sequence of pubertal changes:

- 1) Growth spurt
- 2) Thelarche (Tanner staging I → IV)
- 3) Pubarche (Adrenarche)
- 4) Menarche
- 5) Axillarche

Abnormal puberty



Investigations for all :

- 1) History to detect cause (drug intake)
- 2) US : to detect any tumors
- 3) Hormonal profile : for levels of E₂ , FSH , androgen
- 4) Specific for suspected lesion eg : CT Brain

Menopause (51-52 yrs)

(Reproductive → non reproductive)

Due to depletion of follicles



Clinical manifestations:

- 1) Atrophy of urogenital system :
 - * Vulva : gaping
 - * Vagina : dryness , dyspareunia (ttt : give local E₂ cream) / infection
 - * Cx
 - * Uterus : ET < 5mm
 - * Breast
 - * Urethra / bladder : infection / SUI
 - * Ligaments : POP
- 2) Androgenic manifestations :
 - voice / temporal recession of hair / change in libido
- 3) **HOT FLUSHES (most irritant)** → ttt : HRT
- 4) Mood changes: insomnia
- 5) Skin : loss of collagen
- 6) Remote complications:
 - * CVS manifestations → ++LDL / --HDL
 - * Osteoporosis → ++ osteoclasts / -- osteoblasts

Management:

- 1) Reassurance
- 2) Change life style : exercise / stop smoking / dietary habits
- 3) Prophylactic : Ca⁺⁺ / Vit D
- 4) Annual screening program:
 - TVS / Pap smear / Mammography / Lipid profile / DEXA
- 5) ttt of complications :

- * Osteoporosis
 - Biphosphonate (once weekly)
 - SERM **BUT** ++ hot flushes
 - Phytoestrogen **BUT** weak
 - HRT **BUT** Ca breast / Ca endometrium / ++CVS
 - * **CI : H/O Breast ca / AUB / DVT / Active liver disease**
- Forms :** continuous E/P , Cyclic E/P , E only in TAH
- * CVS complications : prevention / cardiology referral




Ovulatory disorders

{Anovulation}


Type I: Hypothalamic



Type II: Ant. Pituitary



Type III: Ovarian



Type IV: PCO

Cl.picture:

- S : → Regular cycles / spasmodic dysmenorrhea / PMS
- Midcycle → Pain (Mittelschmerz)
- Spotting
- Discharge

Inv :

- 1) BBT chart
- 2) Folliculometry: at midcycle
- 3) PRG: at D₂₁ > 5ng/ml (5-10 = LPD)
- 4) LH Kits urinary : at midcycle
- 5) PEB: not done

Ttt:

- * HMG/ HCG in type I , II
- * ttt of the cause (if possible) in type III
- * PCO (type IV): induction of ovulation

PCO

* Stein & Leventhal 1935 (obesity / Hirsutism / Infertility)

* Rotterdam criteria 2012


(anovulation / hyperandrogen / US necklace appearance)

Irregular cycles hirsutism
Infertility

Inc: 5-10%


Et : theory

++ leptin → ++GnRh & ++ LH (maintained)



Inv:

- I) Labs: → (N) E₂
- ++ androgen
- ++ E₁
- ++ LH / (N) FSH
- - - PRG < 5ng /ml
- ++ insulin resistance

II) US: Adam's criteria  Necklace appearance

III) Lap: (not needed for diagnosis) oyster-shell appearance

Ttt:

- 1) Weight reduction / metformin
- 2) Symptomatic ttt:
 - * COPs to regulate cycles
 - * Spironolactone /cimetidine / OCPs for hirsutism
 - * CC for anovulation (++ OHSS in HMG/HCG)
 - * Corticosteroids in resistant PCO
 - * LOD n resistant PCO (rarely used)

Hirsutism

Def : ++ hair growth in androgen dependent areas

- Et :**
- Constitutional (commonest)
 - Ovarian as PCO
 - Adrenal é ++DHEAS
 - Iatrogenic medication

Diagnosis :

- 1) Labs : * DHEAS
- * Total & free testosterone
- 2) US : for pelvis (ovaries & adrenal)

Ttt:

- * OCPs (++SHBG & -- free androgen)
- * Cimetidine (antiandrogen)
- * Cyproterone acetate (Diane 35) (antiandrogen)
- * Spironolactone (--5α reductase)
- * Finasteride (--5α reductase)
- + hair removal
- Ttt of cause if present

Hyperprolactinemia

Def : ++PRL > 29ng ml

Et :

- * Physiological (commonest)
- * Medication side effect (eg:antipsychotic)
- * Prolactinoma → micro < 1cm
- macro > 1cm
- * Hypothyroidism (↑ TRH)

Diagnosis :

- * Lab: PRL level
- * CT Brain

Ttt : of cause

- * Micro: Bromocriptine (dopamine agonist)
- Or Cabergoline
- * Macro: surgical removal

Infertility

Etiology :

- * Male 40%
- * Sexual 5%
- * Unexplained 15%
- * Female 40%
 - Ovarian
 - Tubal
 - Uterine
 - Cx / vaginal

Cl picture (history):

- **Male :** surgery / drugs / infection...
- **Female:**
 - Ov : ovulation symptoms
 - Tube : operation / PID
 - Ut : amount of menses
 - Cx , vagina : discharge
- **Sexual :** frequency / vaginismus / impotence ...

Inv: → **Male :** semen analysis (± testicular biopsy in azospermic)

* **Count :** > 15 million / ml
 * **Motility:** > 50 % active
 * **Morphology:** > 30% normal
 * **Volume:** > 2ml
 * **PH:** Alkaline

→ **Female :**

- Ov : ovulation tests
- Tubes : HSG
- Uterus : US / HSG

→ **Unexplained :** DHL (diagnostic laparoscopy / hysteroscopy)

Ttt :

- **Male :** stop smoking / control DM / Vitamins / IUI (Intra Uterine Insemination)
- **Female :**
 - Ov : induction of ov (clomid / HMG-HCG)
 - Tubes : IVF (± tubal disconnection in hydrosalpinx)
 - Uterus : ttt of cause eg hysteroscopic myomectomy
- **Unexplained :** IVF-ICSI


IVF-ICSI

Indications:

- Severe tuboperitoneal factor
- Severe male factor
- Unexplained infertility

Steps:

- 1) Pituitary downregulation by GnRha
- 2) Induction of ovulation by HMG



- 3) Final maturation by HCG
- 4) Oocyte retrieval under anesthesia (after 36 hrs)
- 5) Start luteal phase support by natural PRG
- 6) Do IVF-ICSI & observe embryo division , then transfer (D 3-5)
- 7) Continue PRG support for 2 weeks or till end of 1st trimester (if pregnancy occurs)


Factors affecting success rate :

- Maternal age
- Quality of oocytes
- Quality of embryos
- Type of protocol used

Complications : OHSS

- G I , II : mild
- G III , IV: severe + hospitalization

Female sexual response cycle




FSD:
(disorders)
6 categories

- 5) Dyspareunia: persistent pain
6) Vaginismus: persistent involuntary ms spasm

Ttt: For 1) Flibanserin / psychoeducation
For 2) and 3) Testosterone , Tibolone (SERM) , Sildenafil
For 4) Psychoeducation about anatomy
For 5) and 6) Topical lidocaine , antidepressants , pelvic floor physical exercises

Female Genital Mutilation (FGM)

4 types with early (bleeding , infection) & late (loss of satisfaction, sexual problems & frigidity) complications.




Type 1
Removal of clitoris



Type 2
Removal of clitoris
& labia minora



Type 3
Removal of clitoris, labia minora
& labia majora



Type 4
Burning or pricking
or piercing

Fibroid (myoma , leiomyoma)

HMB

- **Definition :** Benign tumour of smooth muscle fibers of the myometrium.
- **Incidence :** 25 % of ♀ in child bearing period !!! WOW 😱
- **Etiology :** ++ E₂ eg (Genetic / Racial / Nulligravida / Anovulation / Early menarche & Late menopause)
- **Pathology :**
 - Gross (site / size / shape / consistency / cut section 🌀 / count)
 - Mic: < 10 mitotic figures / 10 HPF
 - Pathological changes

(Uterine Sarcoma):

- * Rapid growth
- * Rapid recurrence
- * Growth after menopause

Atrophy
Necrosis
Infection


Degeneration →
Malignant (very rare)

- * Hyaline (commonest)
- * Cystic
- * Fatty
- * Calcification (menopause)
- * Red (Necro-biosis) commonest in pregnancy: incomplete necrosis

- **Clinical picture :**
 - S: Asymptomatic / Menorrhagia , unless SM fibroid polyp metrorrhagia
 - Pain / Mass / Infertility...
 - S: General → anemia
 - Abdominal → * symmetrically enlarged ut (submucous myoma)
* asymmetrically enlarged ut (subserous / multiple fibroid)
 - PV & Bimanual → mass (in weeks → describe)

- **Diagnosis :**
 - US: TVS (TAS) : Gold standard
 - Hysteroscopy / Laparoscopy / HSG / MRI / CT / X-Ray...

- **Treatment :**
 - **Conservative:** NO symptoms ✱ NO ttt
EXCEPT : > 14wks / BLM / Infertility or RPL in SMM /
Rapid recurrence / Growth after menopause
 - **Medical (for bleeding / anemia):**
 - Antifibrinolytic eg Tranexamic acid
 - Venotonics eg Daflon
 - **Hormonal (to combat E₂):** Gestagens / GnRh agonist / OCPs
 - **Microinvasive techniques :** UAE / lap. Myolysis / HIFU
 - **Surgical (definitive ttt) :**
 - Myomectomy (open / laparoscopic / hysteroscopic)
 - Hysterectomy (open / laparoscopic / vaginal ?)



NB: Fibroid during pregnancy , Never to be removed , due to high vascularity & ++ E₂ EXCEPT in certain conditions

Endometriosis

PAIN

Def: presence of end. glands & stroma outside ut. cavity

♣ uterus is clean

Incidence: 10% of all ! 🤯 20% of ch. pelvic pain 30% of infertility

NB : Commonest cause of chronic pelvic pain is Idiopathic

Etiology: ++ E₂ / white races

- + theories
- Sampson Retrograde menstruation
 - Halban Lymphatic theory
 - Meyer's coelomic metaplasia theory
 - Immunologic & genetic theory

Pathology:

- site
 - pelvic ✓✓
 - extrapelvic
- size
 - small spots : burn match
 - large cysts : endometrioma (chocolate cyst)

Clinical picture :

Type of patient:

→ **Symptoms:** PAIN / ± Infertility
dysmenorrhea / dyspareunia (deep) / dyschezia / dysuria / ch. pelvic pain

→ **Signs :**

- General
- Abdominal
- Local: PV & Bimanual
 - fixed RVF
 - endometrioma (adnexal mass)
 - nodules in DP

DD of nodules in DP : Endometriosis / TB / Krükenberg

Inv:

- US : for endometrioma (Ground glass appearance)
- CA125: prognostic & follow up not diagnostic
- Laparoscopy : gold standard

Ttt:

- Symptomatic for pain
- Hormonal to atrophy end. glands : **continuous** COPs / gestagens
- Surgical:
 - conservative
 - lap ov. cystectomy if endometrioma > 4cm
 - lap. ablation of end. foci
 - definitive : TAH & BSO

Adenomyosis

HMB

Def: presence of end. glands inside myometrium

♣ uterus is affected

Etiology + theories: more in Multipara
(infiltration in myometrium during involution)

Pathology :

- localized: DD Fibroid (false capsule)
- diffuse

cut section : Granular trabecular pattern with no capsule

Clinical picture :

Type of patient:

→ **Symptoms:**

- bleeding (menorrhagia)
- pain (2^{ry} dysmenorrhea)

→ **Signs:**

- General : anemia
- Abdominal : ± enlarged uterus
- PV & Bimanual:
 - Halban sign (symmetrically tender enlarged uterus)
 - Free adnexae

→ **DD:** symmetrically enlarged ut

Inv: US (TVS / TAS if large Ut)


Ttt:

- Symptomatic:
 - Analgesics / antifibrinolytics / venotonics
 - Hormonal : (to ↓ ↓ menses)
eg : OCPs / gestagens / **mirena LNG-IUD**
- Definitive surgical ttt : TAH

(most common) Atrophic (most serious) EH / EC **After menopause – AUB – Before puberty** (postmenopausal bleeding) mostly FB introduction may be precocious puberty

Child bearing period

(HMB (Menorrhagia) / IMB (Metrorrhagia) / Contact bleeding)



- Polyp
 - Adenomyosis
 - Leiomyoma
 - Malignancy
- (US)
- cx
 - endometrium
 - functioning ovarian tumor

ttt of cause

- Coagulation disorder
- Ovarian disorder (sp after puberty & before menopause/ ...)
- Endometrial disorder
- Iatrogenic drugs (eg anticoagulants)
- Non specific (HTN / Liver troubles / Thyroid troubles)

Ttt:

- of cause if present
- If ovarian disorder:
 - Dysfunctional menorrhagia
 - (mefenamic acid / antifibrinolytic / venotonics)
 - Dysfunctional metrorrhagia
 - COPs / gestagens
 - Mirena IUD (old / multipara)
 - D&C (severe bleeding)
 - End. ablation (alternative to TAH , less morbidity)
 - TAH

Dysmenorrhea

Def :

Types:

- 1^{ry} spasmodic
- 2^{ry} congestive
- Endometriosis (crescendo-decrescendo)

Cl picture:

- Type of pt
- S → type of pain
- Signs → associated symptoms
- Signs → General
- Signs → Abd
- Signs → PV / PR / Bimanual

Inv :

to confirm or exclude organic lesions

Ttt:

- Spasmodic (NSAID / COPs)
- Congestive (ttt of cause)

PMS

(Premenstrual Tension Syndrome)

Def:

Theories :

Inv : Keep diary \$ monthly

Ttt : symptomatic / -- ovulation selective serotonin reuptake inhibitors

***NB :** Metropathia hemorrhagica: Abnormal excessive , often continuous bleeding due to persistence of proliferative phase of menstrual cycle = swiss cheese appearance (back to back arrangement of glands)

Functional ovarian cysts

Characteristics:

- Occurs in childbearing period
- Unilocular
- ≤ 7 cm
- Represents 25% of all adnexal masses
- Ecolucent
- Spontaneous regression is a rule

Types:

- **Follicular cyst (commonest):**
Occurs in 1st half of cycle
- **Corpus luteum cyst (2nd most common):**
Occurs in 2nd half of cycle , vascular & may cause acute pain
- **Theca lutein cysts:**
as in V.mole & OHSS
- **Endometriomas:**
if more than 4cm , laparoscopic cystectomy is needed.
Otherwise medical ttt by continuous COPs
- **Inflammatory cysts (TOC / TOA):**
ttt broad spectrum antibiotics & follow up for size by US ,
if no regression , then surgical drainage is needed
- **Inclusion cysts:**
precursor of epithelial ovarian tumors

ttt: Spontaneous regression is the rule , if not , give cyclical COCPs monthly with US follow up

Pruritis vulvae

- * With vaginal discharge
- * Without vaginal discharge

Etiology:

- 1) dermatoses
- 2) irritation by new medication, soap , condom , toilet paper , ...

Vulval dermatoses

1) Lichen simplex chronicus:

Et :chronic scratching : skin lichenification (beyond labia majora)

Ttt: remove irritant / topical steroid / cool sitz bath / oral antihistamine

2) Lichen sclerosus: usually in post menopausal

Et : may be genetic

CP: thin , pale sp labia minora (hour glass distribution)

Ttt: corticosteroid ointment , if persistent + biopsy

3) Hidradenitis suppurativa (it affects sweat glands in the axilla & mons pubis & causes very bad odor)

ttt : check for DM & local antibiotic cream

4) Crohn's disease:

CP: typical granulomatous linear knife-cut ulcers in labiocural folds & perineum

Ttt: of Crohn's disease oral metronidazol ± corticosteroids

Recurrence is common

Vaginal conditions

- 1) DES induced reproductive tract abnormalities
- 2) Gartner duct cyst

Solid vulval swellings

Malignant : 1ry or metastatic

Benign : condyloma / leiomyoma / fibroma / lipoma

Vulvodynia

Def :

vulvar pain for at least 3 months duration without clear identifiable cause.

Classification :

- * é specific disorder (trauma / infl /neoplastic / hormonal / neurologic)
- * é potential associated factors : localized / provoked

Ttt: individualized

Topical lidocaine , oral antidepressant or anticonvulsant may be used.

Cystic vulval swellings

1) Bartholin gland duct

- * Cyst ttt : marsupialization
- * Infected cyst ttt : antibiotics
- * Abscess ttt: drainage

2) Urethral diverticulum & skene gland

3) Epidermal inclusion cyst : as clitoridal cyst

4) Vulvar hematoma



Contraception

Top Tier: longest duration eg IUD , Sterilization

2nd Tier: hormonal

3rd Tier: barriers & physiological

4th Tier: spermicidal

	Mainly used					In special situation			
Types	IUD	Hormonal					Physiological	Barrier	Sterilization
	* Copper * Silver * Mirena (LNG – IUD)	↓ Oral E / P (daily) P (daily non-stop)	↓ IM E / P (monthly) P (3months)	↓ subdermal Implants (3 years) P	↓ Vag ring E / P (3wks)	↓ Patches E / P (weekly for 3wks)	* safe period * coitus interruptus * lactation sp in first 6 months	*Physical ↳ Condom ♂ / ♀ ↳ Vag diaphragm ↳ Cx cap *Chemical spermicidals ↳ Nonoxynol-9	* Female ♀ Tubal ligation (lap /open / vag) * Male ♂ Bilat vasectomy
Mode of action	* - - implantation (create unfavorable medium ie inflammatory)	* - - ovulation * Atrophic endometrium * Thick cx mucous (Hostile) * Affect tubal motility PRG					* - - Fertilization * - - Ovulation	* - - Fertilization * Kill sperms	- - Fertilization
Benefit	Easy / Cheap / Available								
		Effective						* STDs protection with condom	* Permanent ?
Side effects	(commonly used in developing countries)	* Regular cycles (with COCPs) (commonly used in developed countries)							
	* Bleeding(menorrhagia) except é Mirena * Infection * Perforation / expulsion * Missed IUD * Pregnancy on Top	* Break through bleeding / ↓amount of menses * PMS like \$ * Hypercoagulable state with E containing COPs * Render hypertension , DM difficult to be controlled					* Ineffective * Need cultured couple		* Post ligation syndrome * Permanent
CI	* Infection * undiagnosed bleeding * anatomical ut defect * H/O of: PID/ Ectopic	* H/O of DVT (with estrogen content) * Breast lesion or family H/O * Impaired liver function * HTN / DM * Migraine * Lack of compliance (é pills)					* Irregular cycles	* lack of compliance	* Future desire for fertility (Undecided couple)

* **Emergency contraception:** OCPs : 4 tab $\xrightarrow{12hrs}$ 4 tab , LNG : 0.75 mg $\xrightarrow{12hrs}$ 0.75mg (up to 48hrs) / IUD : (up to 5 days)

* **Contraception during lactation:** All EXCEPT : E content (as Estrogen - - inhibits lactation) / Physiological safe period

Pelvic organ prolapse (POP)

Def : descent ... below (N) anatomical position **Inc :** 25% !!

Et : Weakness of support :

(**lig:** Mackenrodt's / uterosacral / pubocervical / **ms:** levator ani / **fascia**)

due to :
 { Repeated childbirth
 Menopausal atrophy
 Congenital weakness } + PPT factors


Types :

* **Vaginal**
 { ant (urethrocele / cystocele)
 post (deficient perineum / rectocele / enterocele or hernia of DP)
 apical
 (vault prolapse after hysterectomy)

* **Uterine**
 { 1st
 2nd
 3rd / complete procidentia

Pathology :
 { Vagina : keratinization
 Cx : trophic (decubitus) ulcers
 Supravaginal elongation of Cx (sound)
 Bladder : stasis of urine / kinking of ureter

* **Mixed**
 Utero-vaginal
 or
 Vagino-uterine



Cl picture:

* **Symptoms:**

{ Asymptomatic
 Heaviness at end of day / backache / mass protruding
 Symptoms of pelvic congestion
 Ant: urinary symptoms
 (frequency or inability to complete except by reposition)
 Post: rectal (defecation symptoms)
 Sexual symptoms & lack of satisfaction

* **Signs :**

{ Gen : obesity/ chest / abd enlargement
 PV : inspect / palpate / gurgling in case of enterocele
 sound in supravaginal elongation of Cx

Inv :

* Urine analysis (for urinary symptoms)

* Tests for SUI (in cases of cystocele)

Ttt :

* **Prevention :**

{ Proper spacing
 Proper management of 1st , 2nd & 3rd stages of labor
 Kegel's exercise after delivery

* **Actual ttt:** (surgical at least 3-6 months after delivery)

Vaginal:

{ Cystocele : Ant repair
Rectocele : Post repair
Vault prolapse : Abdominal sacrocolpopexy } Cystorectocele:
 classical repair

Uterine :

{ Mild 1st degree : NO ttt
2nd degree : Sacrospinous fixation
2nd degree with cx elongation : Fothergill's or Manchester op.
 (amputation of cx + shortening of Mackenrodt's lig.
 + classical repair)
3rd degree (é need to preserve fertility) : Sacrospinous fixation
3rd degree (é no need to preserve fertility) :
 Vaginal hysterectomy & Classical repair

NB :

* **LeFort operation** : in very old é no sexual activity & can't withstand vag hysterectomy

* **Pessary** : in extremely fragile debilitating , can't withstand any form of anesthesia & surgery

Urinary incontinence (involuntary escape of urine upon ...)

Rectovaginal fistula

	Urethral		Extraurethral
	SUI (urodynamic incontinence)	DO (urge incontinence)	
	Mixed incontinence		GUF → VVF (no desire) → Uretero-vaginal fistula (é desire)
Def	Involuntary escape of urine upon ↑ IAP	Involuntary escape of urine upon desire to micturate	Connection between urinary & genital tracts (Always wet except in urethrovaginal)
Et	* Same as POP * Hypermobility urethra	++ D.contractions (idiopathic / irritation / infection)	* Obstetric (developing countries) * Surgical (developed countries)
Cl.pict	Involuntary escape upon straining	Can't make it to toilet	Always wet * No desire in VVF (except if very small & high) * There is a desire in uretero vaginal from normal ureter
Inv	* Cough test * Bonney's test (in POP +SUI) * Q – tip test > 30° mobility (N) urodynamics : (D.filling pre <15 cmH ₂ O / 1 st Desire 150 ml / Residual < 50ml / full at 400-600 ml) * Normal filling pressure * Leak on cough IVP > IUP	* urine analysis * urine C/S * Cystoscopy * Filling pr > 15cmH ₂ O (due to contraction of detrusor) * No leakage upon cough (as IUP > IVP)	* Sim's speculum & position * Methylene Blue (3 gauze test) * IVP (if KFTs (N)) * Cystoscopy (to visualize fistula)
ttt	SURGICAL Start by - Kegel's - Scheduled voiding * Periurethral plication (Kelly's suture) é 60-70% success * TOT > TVT vaginal sling op (if without POP) * Retropubic urethropexy (Burch colposuspension) (Gold standard) success > 90% * Periurethral injection of Collagen in hypermobile urethra In mixed incontinence: give medical ttt for <u>urge</u> incontinence <u>first</u> , before planning surgery for SUI DD: * Retention with overflow (intermittent self catheterization) * Nocturnal enuresis (psychological ttt)	MEDICAL : bladder training * ttt of infection * ttt of stones * Anticholinergic as oxybutynin (destrusitol)	CATHETER IF SMALL SURGICAL IF LARGE * Dedoublement vaginal in low & abd in high fistulae NB : <u>In uretrovag :</u> desire is present from (N) ureter filling & continuous leakage from affected ureter at site of UVF (– ve MB test / Abd repair) <u>Before surgical repair in vesico vaginal (VVF):</u> * 3-6 months from procedure while putting a urinary catheter to divert urine, decrease size of fistula & allow max healing * Post op: catheter for 10-14 days * No pregnancy for 1 year * Delivery by CS <u>In urethrovaginal:</u> splinting urethra (no incontinence) <u>Vesicouterine:</u> Menouria

Def :
track between rectum & vagina

Cl.pict :
* if small :
incontinence to flatus
* If large:
incontinence to stools

ET:
* mostly between lower rectum & vagina , due to failed episiotomy repair or obstetric trauma.

Ttt: SURGICAL
Preoperative : fluid diet & intestinal antiseptic for 3-5 days
OP :
* **Low1/3 :** transform to complete perineal tear & repair by Lawson Tait operation
* **High fistula:**
Abdominal repair by dedoublement as VVF
Post operative :
* Low residue diet for 7 days
* Use laxatives to avoid constipation
* Episiotomy should be done in subsequent deliveries

Lower Genital Tract Infections (no fever)

Natural barriers ***Vulva:** apposition / apocrine glands é acidic secretions

***Cx:** mucus plug

For LGT

***Vagina:** apposition / st sq ep / E₂ ++ glycogen , lactobacilli act on glycogen → acidic PH (lactic a)

***Ut:** monthly shedding of endometrium

LGT infections:

Vulvitis / Vaginitis

Cervicitis

Childhood: foreign body / worms as oxyurias

Menopause : -- E₂ Atrophic / Alkaline PH

Childbearing

vaginitis

	Bacterial vaginosis (commonest)	Candidiasis (2 nd common)	Trichomoniasis (3 rd common)
organism	Bacteria Gardnerella vaginalis	Fungus Candida albicans	Protozoal Trichomonas vaginalis
Et	++++ anaerobes	--- immunity	STD
PH	Alkaline : 4.7-7	Acidic < 4.5	Alkaline : 5-6
Cl.picture	Fishy odour Profuse / non irritant Grayish discharge	Extremely irritant Cottage cheese Scanty Odorless discharge	Profuse / yellowish Mildly irritant / malodorous discharge / é BV & other STDs Strawberry Cx In 25% of cases
Mic +KOH	Clue cells 	Hyphae / pseudohyphae	Flagellated protozoon 
Treatment	Metronidazole 500 mg bid / 7days (CI in 1 st trimester, give local cream or supp only)	Fluconazole 150 mg once weekly for 2 weeks (CI all through pregnancy & in liver troubles, give local cream or supp only)	Metronidazole 2gm single oral dose + ttt of partner

Cervicitis

	Acute	Chronic	Erosion
Etiology	Polymicrobial sp. Neisseria & Chlamydia	Recurrent improperly ttt acute	*infection (ch.cervicitis) *hormonal(preg /COCPs) *congenital
Cl.picture	*Mucopurulent discharge *Dyspareunia *Backache	*Leucorrhea due to pelvic congestion *Mucous polyp *Nabothian follicles *Hypertrophic Cervicitis	Contact bleeding
Inv.	We may do C/S from discharge in cases resistant to ttt	—	Pap smear
Treatment	Doxycycline 100 mg bid / 7days Or Azithromycin 1gm single oral dose (in pregnant females)	*ttt of infection in case of acute on top of chronic *ttt of erosion in case of erosion	* ttt of the cause. * Residual cases: Cauterization → Diathermy → Cryo → Laser

Upper Genital Tract Infections

Acute PID (presence of fever)

Definition: Salpingitis , oophoritis , peritonitis & rarely endometritis due to regular monthly shedding in childbearing period.

- Et:**
- **organism:** N.gonorrhea, Ch.Trachomatis (mostly both together& others)
 - **route:** ascending (mainly), local or lymphatic, blood (rarely)
 - **risk F:**
 - sexually active , multiple sexual partners
 - IUD , after menses , after sexual intercourse
 - é any procedure : D&C , HSG , Hysteroscopy, ...

NB:

- OCPs & barrier contraception ↓ risk (ie are protective)
- Chlamydia may cause silent PID

Pathology:

- Endosalpingitis / interstitial salpingitis / perisalpingitis (acute catarrhal or acute suppurative)
- Oophoritis : é microabscesses on the surface
- Pelvic peritonitis

Clinical picture:

- **Symptoms:**
 - fever , malaise , headache ,
 - H/O of recent OBGYN procedure
 - Acute lower abdominal pain
 - Foul smelling purulent vag. discharge
- **Signs:**
 - Fever $> 38.3^{\circ}$, tachycardia
 - Lower abdominal tenderness & rebound tenderness
 - Cx Motion tenderness , discharge
- **Complications :**
 - Abscess / septicemia
 - Chronic PID (if inadequately treated)
 - Tubal obstruction & infertility (due to fibrosis & adhesions)
 - ++ risk of ectopic pregnancy
- **DD:** acute abdomen

Inv:

- **Blood tests:** ++TLC , ++DLC , shift to Lt , ++ESR , ++ CRP
- **US** to exclude DD : ectopic , complicated ovarian , appendicular mass , degenerated myoma
- **Laparoscopy:** if diagnosis is doubtful

NB:

- Exam of discharge to detect causative organism & do C/S (not clinically needed)
- Fitz Hugh Curtis syndrome (filmy adhesions between the liver & under surface of diaphragm may be seen on laparoscopy in cases of chlamydia)

Treatment:

- **Mild PID (é mild symptoms) → Outpatient**
 - Ceftriaxone 250mg single IM dose
 - + doxycycline 100mg oral /12hrs/14days
 - ± Metronidazole 500mg oral bid / 14days
- **Severe PID (é severe symptoms) → Hospitalization**
 - IV fluids /IV analgesics / IV antipyretics
 - IV antibiotics :
 - Cefotetan 2g IV/12hrs or cefoxitine 2gm IV/6hrs
 - ⊕ Doxycycline 100 mg orally / 12hrs
 - till symptoms become milder then continue previous oral regimen of mild cases for 2weeks

NB:

- 1) IF IUD present → remove
- 2) In case of TOA → give clindamycin 900mg IV / 8hrs or Metronidazole 500 mg IV / 8hrs in addition to the 2 above mentioned drugs of severe PID
- 3) IF TOA doesn't resolve by medical ttt :
 - drainage through laparoscopy / laparotomy or colpotomy (vaginal drainage)

Chronic PID

Sequelae of acute

Pathology:

- Hydrosalpinx : sequelae of inadequately treated acute catarrhal salpingitis
- Pyosalpinx : sequelae of inadequately treated acute suppurative salpingitis
- TOC
- TOA
- Chronic Interstitial salpingitis

Clinical picture :

Symptoms :

- * history of acute PID
 - * dull aching lower abdominal pain
 - * pelvic congestion
 - * backache
 - * infertility from tubal obstruction & peritoneal adhesions
- Menorrhagia
Leucorrhea  Congestive dysmenorrhea + Dyspareunia

Signs:

- * adnexal tenderness/ fullness / cyst
- * fixed RVF in case of extensive adhesions

investigations:

- US for adnexal masses
- HSG for hydrosalpinx
- Laparoscopy is the gold standard in cases of ch. pelvic pain

Treatment :

Symptomatic ttt

- eg : → infertility ttt → (TD + IVF)
 → pain & congestion in old age → (TAH & BSO)
 (antibiotics only in acute exacerbations)

TB (chronic from the start)

TB is a chronic granulomatous disease

Etiology:

- Mycobacterium tuberculosis
- Blood spread from lungs (most common)

Pathology : (genital TB)

- FT: (affected in 100% of cases of TB of genital tract)
- * **Endosalpingitis**: caseous material inside thick , tortuous , tobacco pouch appearance (open everted fimbrial end)
- * **Interstitial salpingitis** : thick , beaded , salpingitis ithmica nodosa
- * **Perisalpingitis**: é multiple tubercles on the surface & on surrounding peritoneum
- Endometrium affection (50 %) : affection of basal layer → IU adhesions or Asherman syndrome (PEB is diagnostic)
- Ovarian affection (25%) : with microtubercles
- Rarely : cervical, vaginal or vulval serpiginous ulcers é undermined edges & necrotic floor

Clinical picture :

Symptoms

- Asymptomatic / ch.pelvic pain
- Of 1^{ry} / Low grade fever/ loss of wt / loss of appetite
- 5-10% of infertility cases are due to TB salpingitis
- Amenorrhea / oligomenorrhea / hypomenorrhea

Signs:

- Mostly normal + Nodules in DP
- Genital serpiginous ulcers é undermined edges

Investigations :

- X-ray chest (& pelvis for calcified LNs)
- HSG (not in active TB) : retort shape tube , IU adhesions
- PEB
- Laparoscopy ± biopsies from suspicious lesion (serosal tubercles to be stained by Ziel –Nielsen to show the acid fast , alcohol fast bacilli) & cultured in Löwenstein-Jensen medium
- Tuberculin is a good –ve test

Ttt:

- General ttt for anemia , proper nutrition
- Anti TB (Rifampicin , INH , Pyrazinamide , Ethambutol)
- Surgical ttt in case of tubal mass or endometrial TB

STDs

	Bacterial					Viral			\$
	⚡	Chlamydia Trachomatis	Chancroid	Donovanosis (GI)	LGV	HSV ₂	HPV	HIV	
Organism	Neisseria gonorrhea (columnar & transitional)	Chlamydia Trachomatis (columnar epithelium)	Hemophilus Ducreyi	Klebsiella granulomatis	Chlamydia L _{1,2,3}	HSV ₂	HPV 6 / 11	AIDS	Treponema Pallidum
Cl.picture	PID		Vulval lesions					Vulval + Systemic manifestations	
	- Mucopurulent discharge - Lower abdominal pain - Bartholin - Urethritis - Systemic (IP 3-5 days)	- Subclinical Mucopurulent discharge - PID (silent) - Squelae: Fitz Hugh Curtis syndrome in laparoscopy	<u>Painful</u> papule ↓ painful ulcer é exudation LN+++ (IP 3-5 days)	<u>Painless</u> papule ↓ Ulcerate ××× NO ××× ××× LN××× (IP 3 weeks)	Destructive lesion ↓ Ulcerate LN+++ (IP 3weeks)	<u>Painful</u> Vesicles ↓ painful ulcer without exudation (IP 3weeks)	<u>Painless</u> warts (condyloma acuminata)  (IP 3months)	Asymptomatic ↓ Severe form Kaposi sarcoma (IP 3years)	1^{ry} (chancre) <u>Painless</u> 2^{ry}(condyloma latum) (maculopapular rash) 3^{ry}(systemic) (tabes dorsalis)& gumma formation Congenital: Vertical transmission
Inv.	- Gram -ve diplococci - Culture: Thayer Martin NAAT	- Obligatory intracellular - Culture : expensive NAAT	- Coccobacilli - Cl.picture is enough	Gram -ve Donovan Bodies India/Africa	CF India/Africa	- Culture of serum collected from vesicles - Ab	- Pap smear (koilocytes) - Colposcopy	- Western blot - Eliza	- Dark field Mic (spirochetes) - Non-specific VDRL / RPR - Specific:TPH/TPI - In 1^{ry} \$ ▶ Ag present - In 2^{ry} \$ ▶ Ag & Ab - In 3^{ry} \$ ▶ Ab present
Ttt	Ceftriaxone 250mg IM + Doxycycline 100mg / bid / 7days	Doxycycline 100mg / bid / 7days	Ceftriaxone 250mg IM	Doxycycline 100mg / bid / 3 weeks	Doxycycline 100mg / bid / 3weeks	- Acyclovir 400mg 10 days (can be taken all through pregnancy) - Vesicles at labor = CS	Vaccine - Cryo - Diathermy - Podophyllin - Podofilox - NB : CS only if large lesions obstructing labor	Antiretroviral ttt Vaccine ?	Penicillin In penicillin allergy: - If pregnant: Desensitization - If not pregnant: Doxycycline

NB: in case of pregnancy Doxycycline 100mg / bid / 7 days is substituted by Azithromycin 1 gm single oral dose as doxycycline is teratogenic (but not for syphilis)

Other STDs : Trichomoniasis (protozoon) → infecting lower genital tract / Pediculosis pubis & scabies (Ectoparasites)

Scheme for oncology

Definition: From where the tumor arise

Incidence: How common / rare

Etiology :

- Predisposing F:
- Premalignant lesions :

Pathology:

- Gross :
 - * **Ulcer** with raised everted edges , irregular necrotic floor , indurated base
 - * Cauliflower **mass** with areas of hemorrhage & necrosis
 - * Firm / hard endophytic or exophytic **nodule**
- Microscopy : (depends on cell of origin)
 - eg : squamous cell carcinoma / adenocarcinoma if arising from glands
- Grading :
 - *G I → < 5% malignant undiff.cells = best prognosis
 - *G II → 5-50% malignant undiff.cells = intermediate prognosis
 - *G III → > 50% malignant undiff.cells = poor prognosis
- Spread :
 - * Direct spread to surrounding structures
 - * Lymphatic to the draining LNs
 - * Blood : Lung Bone Liver Brain

Cl . picture :

- Symptoms:
 - related to function / symptoms of metastasis
- Signs:
 - * General: for metastasis
 - cachexia / anemia / jaundice / virchow's LN...
 - * Abdominal:
 - * PV & bimanual :
- Staging:
 - * Stage I : confined to the organ
 - * Stage II : limited local spread
 - * Stage III : more local spread ± LNs
 - * Stage IV: distant spread
 - IV a : mucosa of bladder & / or rectum
 - IV b : distant spread (L B L B)

Inv.

- To confirm diagnosis :
- To detect spread:
 - * eg: chest x-ray / abdominal & pelvic US / bone scan
- To assess fitness of pt for surgery: ECG & Lab testing

Ttt:

- Stage I , II :  Surgery
- Stage III , IV :  Radiotherapy / Palliative

Endometrial carcinoma



Def: Tumor arising from Endometrial glands

Inc: commonest tumour of ♀ genital tract & having best prognosis

Etiology :

- Predisposing F: ++ E₂ (unopposed) → early menarche / PCO
late menopause / Granulosa cell tumor / use ERT / obesity / Tamoxifen
- Premalignant lesions : EH → simple 1% / complex 3%
simple é atypia 9% / complex é atypia 29%

Pathology:

- Gross
 - Localized: endometrial polyp
 - Diffuse: ++ endometrial thickening > 5mm
- Microscopy
 - Adenocarcinoma (best prognosis) & commonest
 - Adenoacanthoma (+ benign sq metaplasia)
 - Adenosquamous (+ malignant sq cells)
 - Clear cell ca / papillary cell ca (undiff. so poorest prognosis)
- Grading :
 - G I → < 5% malignant undiff.cells = best prognosis
 - G II → 5-50% malignant undiff.cells = intermediate prognosis
 - G III → > 50% malignant undiff.cells = poor prognosis
- Spread :
 - Direct: myometrium / Cx / Ovaries / FT / vagina
 - Lymphatic: para aortic / inguinal / paracervical / parametrial
 - Blood : L B L B

Cl . picture :

- Type of pt / History
(postmenopausal / white race / NG / PCO / Tamoxifen)
- Symptoms: Post menopausal bleeding (commonest)
- Signs:
 - *Gen: for metastasis : cachexia/ anemia / jaundice / virchow's LN...
 - *Abd: enlarged soft uterus ± signs of metastasis : ascites / liver
 - *PV & bimanual : enlarged soft uterus
± adnexal masses (? ut to ov or ov to ut)
- DD :of post menopausal bleeding

Staging: (FIGO surgical 2018)

- Stage I : confined to the organ (uterus)
 - I a < ½ myometrial invasion
 - I b > ½ myometrial invasion ± endocervical glands
- Stage II : limited local spread : Cx. stroma
- Stage III : more local spread ± LNs
 - III a : serosa of uterus and / or adnexa
 - III b : Vagina / Parametrium
 - III c : LNs (parametric LNs / paraaortic LNs)
- Stage IV: distant spread
 - IV a : mucosa of bladder & / or rectum
 - IV b : distant spread (L B L B)

Inv.

- To confirm diagnosis :
 - * Screening:TVS (if ET > 5mm in menopausal) / Hysteroscopy
 - * Gold standard (confirmatory) : FC & endometrial biopsy
- To detect spread:
 - * eg: chest x-ray / abdominal & pelvic US / bone scan
 - * MRI for myometrial invasion & for LNs
- To assess fitness of pt for surgery: ECG & Lab testing

Ttt: * Early : Surgery * Late : Radiotherapy

- EH :
 - depends on:
 - type of EH
 - age of pt
 - desire for fertility
 - options:
 - PRG (DMPA / Mirena)
 - Hysterectomy
- Stage I : TAH & BSO + cytology ± later radiotherapy
- Stage II : ttt as cancer Cx (Wertheim's operation)
- Stage III : radiotherapy
 - External beam = pelvis
 - Extended = pelvis + abdomen
- Stage IV : Palliative : pain relief / radiotherapy

NB : Unfit pt for surgery in stage I , II : Radiotherapy

- * **Ut sarcoma** : rare after fibroid / blood spread / ttt: hysterectomy
- * **Choriocarcinoma** :
 - low risk / high risk
 - blood spread

Cancer Cervix



Def: Tumor arising from ectocx (70%), endocx (30%) starting in reserve cells of TZ

Incidence: 2nd most common after endometrial Ca. / **most preventable**

Etiology :

Predisposing F: → Sexuality: early / multiple / low socioeconomic / multipara / smoking
→ Viral: HPV (16 , 18) / HSV₂ / HIV

Premalignant lesions: CIN 1 (LSIL) / CIN 2,3 (HSIL)

CIN 1: atypical cells involve lower $\frac{1}{3}$ of ep.

CIN 2: atypical cells involve lower $\frac{1}{2}$ (or $\frac{2}{3}$) of ep.

CIN 3: atypical cells involve all layers of ep.

without
invasion
of BM

Pathology: → **Gross:** Ulcer , Nodule , Friable mass (ectocx) / Barrel-shaped cx (endocx)

→ **Microscopy:** Sq cell Ca (ectocx 85%) / Adenocarcinoma (endocx 15%)

→ **Grading :** → G I → < 5% malignant undiff.cells (best prognosis)
→ G II → 5-50% malignant undiff.cells (intermediate prognosis)
→ G III → > 50% malignant undiff.cells (worst prognosis)

→ **Spread:** → **Direct:** uterus / vagina / parametrium / uterosacral / bladder / rectum.

→ **Lymphatic:** → 1^{ry} : paracervical / obturator / ext iliac / int iliac
→ 2^{ry} : common iliac / lat. sacral / para aortic

→ **Blood :** L B L B

Cl . picture :

→ **Symptoms** → type of patient / **Contact bleeding**

→ **Signs** → **General:** Signs of metastasis (anemia – jaundice – virchow's LNs...) / **Uremia**

→ **Abdominal:** **N**

→ **PV& bimanual :** Nodule / Mass / Ulcer / vagina / normal sized uterus

→ **PR :** uterosacral & Parametrial involvement

→ **Staging (Clinical FIGO 2018)**

→ **Stage I (confined to cx) :** → I a < 5mm depth of invasion

→ I b > 5mm depth of invasion

→ **Stage II (local spread) :** → II a: vagina (but not lower $\frac{1}{3}$)

→ II b: parametrium (but not to lat pelvic wall)

→ **Stage III (more local ± LNs) :** → III a : vagina (+ lower $\frac{1}{3}$)

→ III b: parametrium till lat pelvic wall (**Uremia & Death**)

→ III c : LNs

→ **Stage IV (distant) :** → IV a: mucosa of bladder & rectum

→ **DD:** Contact bleeding → IV b: distant spread L B L B

Inv.

→ **To confirm diagnosis :**

1) Pap smear :new liquid based technique (screening)

2) Colposcopy 3) HPV test

4) Biopsy (direct / colposcopy /punch / Cone / LEEP)

→ **To detect spread:** CXR / Abd & Pelvic US / Bone scan
Ba enema / EUA / IVP / Cystoscopy / ...

→ **To assess fitness of pt. for surgery :** ECG / Lab tests

ttt

→ 1^{ry} prevention (HPV vaccine for teenage)
only if unaffected by HPV

→ 2^{ry} prevention : screening program by pap smear for
whole population / 3years in low risk ,
annually in high risk

→ **CIN** → 1→ Medical ttt & Pap smear after 3-6 months

If refractory to medical ttt :

Cryocautery / diathermy cautery / laser cautery

→ 2→ LEEP / LLETZ

→ 3→ Conization in young age (to preserve fertility)

→ TAH in old age (not desiring fertility)

→ **Cancer Cx:**

* Stage I a : conization if young

* Stage I b : Trachelectomy + abd cerclage if young

* Stage II a : Wertheim's operation (radical hysterectomy)

* Stage II b / III / IV: Radiotherapy / palliative

NB :

* **Radiotherapy can be used in early stages giving same results as surgical ttt**

* **Unfit patient : Radiotherapy in all stages**

* **Ovaries may not be removed as it is not a hormone dependent cancer**

* **In recurrent cancer Cx : use opposite line of what was used before**


* **Ca. of cx stump**

→ difficult surgery (adhesions)

→ no room for intracavitary radiation

→ better prevention by TAH instead of subtotal hysterectomy

Epithelial 70% ←
Germ cells 20 - 30% ←
Sex cord stromal < 5% ←



Benign ovarian tumors (premalignant ovarian lesions)

Clinical classification: Cystic / Solid
Pathological classification: Benign / Malignant
Histological classification

		Epithelial tumors (post menopausal)				Germ cell tumors (child bearing)				Sex cord stromal			
		(Tubal ep)	(Endo Cx ep)	(End)	(urinary ep)	embryonic (endoderm , meso,ectoderm	extra embryonic villi	undifferentiated germ cells & sex cord	Granulosa cells	Sertoli Leydig	Fibrous tissue	Theca cells	
Gross	Size	Moderate	Huge	Endometrioid (malignant from start)	Brenner	Dermoid (BCT)	Yolk Sac (Endodermal sinus tumor) malignant	Gonadoblastoma	Sertoli-leydig cell tumor (androblastoma)(malignant)	Fibroma	Thecoma		
	Site	bilateral in 50%	unilateral		Small	Moderate		Small		Small-Moderate	Small		
	Consistency	Uni/multilocular Papillary (exo/endophytic)	Multilocular		bilat in 15%	bilat in 15%		bilateral in 15%		bilat 10%	bilat 10%		
Microscopy	Characteristic features	Cuboidal ep (ciliated / Non)	Columnar ep é Goblet cells		Solid	Uniloc é thick capsule & long pedicle		Solid		Solid é long pedicle	Solid		
		CA 125	CA 19-9		Transitional ep	Endo / Ecto / Mesoderm Rokitansky nodule		Undifferentiated Germ cells & Sex cord cells		Fibroblast	Theca interna cells		
Secretions				Coffee bean nuclei	hair/teeth /bone - mamilla - chemical Peritonitis	In dysgenetic gonads Y ch . as AIS	Meig's syndrome (+ ascites & Rt pl.effusion)	Post menopausal					
Malignant transformation		30% cystadenocarcinoma	5%	± E ₂	Struma -ovarui (thyroxine)↗	—	E ₂						
	<td></td> <td></td> <td>—</td> <td>< 1%</td> <td>30% Dysgerminoma</td> <td>—</td> <td>—</td>			—	< 1%	30% Dysgerminoma	—	—					

Complications:

- 1) Torsion → gangrene (rare)
- 2) Hemorrhage → acute abdomen
- 3) Rupture → nothing (in serous)
→ chemical peritonitis (in dermoid)
→ pseudo-myxoma (in mucinous)
- 4) Infection → in puerperium
- 5) Incarceration → pressure symptoms
- 6) Malignant transformation

Cl.picture:

- symptoms:
- * asymptomatic
 - * mass (abdominal)
 - * no bleeding (except functioning)
- signs:
- general : Cachexia in Mucinous
 - abd: swelling (insp. / palp. / percussion)
 - PV & Bimanual : Adnexal mass
 - DD Krükenberg tumor
- * acute abdomen (pain)
* pressure symptoms

Inv : **US** / CA 125 / ± Laparoscopy

Ttt:

- Ovarian cystectomy
 - Oophorectomy (in huge mass)
 - Panhysterectomy (TAH+BSO) (in old age)
 - In pregnancy (when to remove it ?)
- (whether laparoscopic or laparotomy)

Low Malignant potential (LMP) OR Border line ovarian tumors


Having intermediate features between benign & malignant cancer

Unlike benign	Unlike malignant
Have at least 2 of the following - nuclear atypia - epithelial stratification - Mic: papillary projections cellular pleomorphism mitotic activity	Lack stromal invasion , But May have Mic stromal invasion < 3mm

Diagnosis : CA125 , US

Treatment : as malignant

If cystectomy , residual disease is left



Malignant ovarian tumors

Def : malignant T. arising from surface ep , germ cells & sex cord stroma of the ovary.

Inc : 3rd most common malignancy of ♀ genital tract é worst prognosis , deaths from ovarian tumors > deaths from end. & cx cancer together

Et : ➔ **Predisposing F :** NG / induction of ovulation / white race / genetic (BRCA 1 & 2 , Lynch 2) , **OCPs are protective as they - - ovulation**
➔ **Premalignant :** Epithelial T (serous 30% & mucinous 5% cystadenoma) / Germ cells (BCT < 1% / Gonadoblastoma 30%)

Pathology : Gross : bilateral / solid é hemorrhage & necrosis / ascites / vascularity

Microscopic & Grading

	Epithelial (> 70 %) old age			Germ cell Tumors (20 - 30 %) Child bearing				Sex cord stromal Tumors (< 5 %) Extreme ages	Metastatic	
	Serous	Mucinous	Endometrioid	MCT MST	Chorio Ca	Endodermal sinus T	Dysgerminoma	Granulosa cell T	Sertoli Leydig	Krükenberg (atypical 2 ^{ries})
Mic	Psammoma bodies	—	End. glands	Endo / Meso / Ectoderm	Trophoblasts	Shiller Duval bodies	Lymphocytes	Call Exner bodies (Rosette shape)	Androgenic cells	Signet ring cells 
Secretions		CA19-9 CA 125		—	HCG	AFP	LDH	E ₂ / Inhibin	Androgen	—

Spread :

- ➔ **Direct:** other ovary / uterus / fallopian tubes
- ➔ **Lymphatic:** para aortic LNs
- ➔ **Blood :** L B L B
- ➔ **Transcoelomic:** peritoneal seedling

Cl.pict:

- ➔ **Symptoms :** Type of pt / Asymptomatic (very late presentation) / GIT symptoms / mass / cachexia / NO bleeding except in functioning T
- ➔ **Signs:**
 - General: cachexia / anemia / jaundice / virchow's LNs / pl effusion
 - Abd: inspection / palpation / percussion (mass /ascites) peau d'orange
 - PV& Bimanual: Nodules in DP in krukemberg T / Adnexal masses

➔ **Staging:**

- Stage I (confined to ovary) **Ia:** one ovary , **Ib:** both ovaries
Ic: one or both é + ve peritoneal cytology
- Stage II (local spread: adnexa) **II a:** Ut / FT , **II b:** UB / colon / rectum
- Stage III (above pelvic brim ± LNs) **III a:** Mic , **III b:** < 2cm , **III c:** > 2cm
- Stage IV (distant spread) ➔ **IVa:** lung & pleural effusion
➔ **IVb:** distant liver parynchyma / brain / bone

FIGO surgical
staging 2018

Inv :

➔ **To confirm diagnosis**

US: bilateral / solid / papillae /ascites (1 or more)

Doppler for vascularity / Tumor markers (CA 125) / Laparoscopy

RMI = US (1 or 3) × Menopause (1 or 3) × CA 125 ➔ **IF > 200 (High risk)**
(1 if one criterion in US or 3 if more than one criterion in US)

➔ **To detect spread:** Ba meal / enema / upper & lower GI /CXR /CT abd & pelvis

➔ **To assess fitness of pt for surgery :** ECG / Labs /...

Ttt: Early : surgery / Late : surgery (debulking + chemotherapy)

Surgery (in all stages)

Stage I :Exploratory laparotomy and TAH + BSO + omentectomy + peritoneal cytology + sampling LNs

Stages II ➔ IV : Debulking + **chemotherapy** as Carboplatin or Taxol

NB:

- * Debulking in late stages with fixed tumor (as above + removal of all tumor tissue > 1cm) :to decrease toxicity of post operative chemotherapy
- * Stage Ia (germ cell T):can be unilat salpingo oophorectomy till completing her Family
- * Dysgerminoma : radiosensitive

Tumors of the vulva

Def: Malignant tumor arising from labia majora / minora / clitoris .

Inc: 4th most common tumor of ♀ genital tract after end , cx & ovaries

Etiology (VIN) :

- Vulval LSIL : as condyloma acuminata (HPV 6,11)
- Vulval HSIL: with oncogenic HPV 16
- Differentiated VIN : as Lichen sclerosus

Pathology:

- Gross:
 - VIN : may appear **N**
 - Invasive: Nodule / Ulcer / Mass
- Mic: Invasive:
 - Squamous cell ca.(commonest 90%)
 - Paget's Adeno ca. / Melanoma / Basal cell ca (rare)
- Grading
 - G I : < 5% malignant undiff.cells (best prognosis)
 - G II: 5-50% malignant undiff.cells(intermediate prognosis)
 - G III : > 50% malignant undiff.cells (worst prognosis)
- Spread :
 - **Direct:** vagina / urethra / anus / perineum
 - Lymphatic: inguinal → femoral (LNs of Cloquet)
 - Blood : L B L B

Cl . picture :

- Symptoms
 - Type of pt
 - In VIN : Asymptomatic or Pruritis vulvae
 - In invasive : Pain / Mass / Ulcer / Contact bleeding
(in addition to long standing pruritis valvae)
- Signs :
 - *General: signs of metastasis
 - *Abdominal : rare
 - *Local:
 - In VIN :
 - raised rough skin / thin reddish epithelium
 - white hyperkeratinization (leukoplakia)
 - In invasive:
 - ulcerated / pigmented mass
 - labia majora mostly then minora then clitoris
 - ± enlarged inguinal LNs

Staging: (FIGO surgical)

- Stage I : (confined to vulva) : Ia : ≤ 2cm é depth ≤ 1mm
Ib : > 2cm é depth >1mm
- Stage II: (lower urethra , lower vagina)
- Stage III: Stage II + Inguinofemoral LNs
- Stage IV: (distant spread)
 - * IVa: UB mucosa / Rectal mucosa
All urethral mucosa / All vaginal mucosa
 - * IVb: L B L B

Investigations:

- To confirm diagnosis: Direct excisional biopsy
If no apparent lesion: use colposcopy or paint é Toluidine blue & take biopsy from the white lesions
- To detect spread :
 - Colposcopy for cervix & vagina to detect associated involvement
(as all are predisposed by HPV infection)
 - CXR / CT abdomen & pelvis
- To assess fitness of pt. for surgery : ECG & Lab tests

Treatment:

- Vulval LSIL : self limited
- Vulval HSIL : local excision **or** topical 5- flurouracil
CUSA (Cavitational Ultrasonic Surgical Aspiration) in non hairy skin
- Vulval cancer:
 - * wide local excision
 - * simple vulvectomy + sentinel LN biopsy
 - * radical vulvectomy
- Radiotherapy or chemo-radio in unfit patients**

Tumors of the vagina

*Least common

* 2ry far more common than 1ry


ttt : * in VaIN : 5- flurouracil vaginal cream for multifocal lesions




*Surgery : Wertheim for early cases



* Radiotherapy for most of cases

NB: Sarcoma Botryoides in prepubertal girls → Mass & Bleeding

Anatomy of female genital tract

	Gross	Histology	Arterial supply	Lymphatics	Applied anatomy
Vulva	 - Mons venereum (hairy pad of fat over SP) - Clitoris: midline erectile organ - 2 labia majora: lat hairy é fat , sweat & sebaceous glands - 2 labia minora: med é non hairy skin , no fat - Hymen : memb 2 cm deep to vulval opening - Bartholin glands: on posterolat aspect of labia majora , duct opening in vestibule - Skene's tubules: paraurethral glands	- Stratified squamous epithelium (keratinized) - Transitional ep. For Bartholin gland	- Int pudendal artery (one of 2 terminal branches of ant division of IIA)	Superficial inguinal LNs	- Clitoridal cyst : post circumcision - Cryptomenorrhea:in imperforate hymen - Bartholin cyst : in obstructed duct
Vagina	- Fibromuscular potential tube é rugae - 7 – 9 cm anterior wall related to UB & urethra - 10 – 11 cm posterior wall related to DP , rectum , perineal body - Lat borders related to : 1) ureter 1cm 2) cardinal (Mackenrodt's) lig 3) levator ani (deep perineal ms) 4) bulbocavernous (superficial perineal ms)	- Non keratinized stratified squamous epithelium	- Vaginal artery from IIA	Upper third : as Cx Lower third : to Inguinal LNs Middle third : II LNs	- Septum : longitudinal / transverse - Ant. wall prolapse: Cystocele , urethrocele - Post wall prolapse: Enterocoele , rectocele , deficient perineum - Support : Mackenrodt's lig / Uterosacral lig / Pubocervical lig - Episiotomy : cut through post or posterolat vag wall - Post colpotomy or culdocentesis : for drainage of pelvic abscess in DP - US guided oocyte retrieval through post colpotomy - Ureter may be injured at lat fornix while clamping the vag angles in TAH - Pudendal n block :done at level of ischial Spines

	Gross	Histology	Arterial supply	Lymphatics	Applied anatomy
Uterus	- Pear shape 1× 2× 3 inches  - AVF (85%) or RVF (15 %) - Ant : UB / Pubocervical lig - Post : DP / Uterosacral lig - Lat: Broad lig & inside it : - Fallopian tubes - Remnants of Wolffian ducts - Uterine artery - Ureter & Mackenrodt's lig	- Endometrium (modified mucosa é glands & stroma) - Myometrium of body of ut formed of 3 ms layers : - Outer longitudinal - Middle criss-cross - Inner circular While Cx is formed of outer & inner layers only - Perimetrium : peritoneal adherent over the body but loose over Cx	- Uterine a (tortuous course on lat borders) from ant division of IIA - Anastomosis at cornu with ovarian a	- Fundus : paraortic LNs - Cornu : superficial inguinal LNs - Body : II LNs - Cx : 1^{ty} → paracervical parametrial , obturator , II LNs & EILNs 2^{ty} → CI LNs , Lat. sacral & Paraortic LNs	- Ut prolapse : 1st , 2nd & 3rd degrees - Ut. support : Mackenrodt's lig / Uterosacral lig / Pubocervical lig (as vagina) - MRKH Syndrome in case of 1^{ty} amenorrhea é presence of secondary sexual characters - Cong anomalies : as septate / bicornuate / didelphys with RPL , PTL - Cx is sensitive only to dilatation (Cx dilatation should be done under anesthesia)
Fallopian tubes	2 Tortuous tubes 10cm in length  - Runs in upper part of broad lig - For ovum pick up , transport & nutrition for ovum & sperm	- Mucosa (endosalpinx) cubical or columnar partially ciliated - Musculosa: 2 ms layers (outer long & inner circular) - Serosa or loose peritoneal covering	- Branches from uterine & ovarian arteries	Paraortic LNs	- Tubal point (½ inch above mid inguinal point) in cases of pain , PID , ectopic . - Commonest site for ectopic pregnancy (ampullary part) - Hydrosalpinx & pyosalpinx (ch . PID) - Commonest permanent method of contraception (tubal ligation) - Tubal disconnection (TD) in cases of hydrosalpinx prior to IVF
Ovaries	- Almond shape 1×2×3cm - On ureter & bifurcation of II vessels - Not covered by peritoneum - Corrugated surface due to repeated stigmata of ovulation - Attached to ut by ovarian ligament , to lat pelvic wall by infundibulopelvic lig , to the back of the BL by mesovarium	 1) Outer cortex é follicles 2) Inner medulla 3) Hilum (site of attachment of mesovarium)	- Ovarian arteries from abdominal aorta NB : - Rt ov. v → IVC - Lt ov. v → Lt Renal v	Paraortic LNs	- In ovariectomy : 6 clamps needed 2 on each pedicle (ovarian , infundibulopelvic & mesovarium) ligaments. - Streak gonads in absence of sex cord cells covering the germ cells , forming Turner syndrome (45 XO) - Responsible for 1) ova production (ie ovulation) 2) hormone formation (E₂ , PRG) by granulosa & theca cells of oocytes

Ureter	Pelvic floor	Perineum
Gross : 25 cm in length , retroperitoneal - Enters pelvis in ovarian fossa above the bifurcation of CIA - Runs downwards in the base of BL <u>below</u> uterine artery . - crosses forward related to lateral vaginal wall to enter trigone of the urinary bladder. Histology : Lined by transitional epithelium Arterial supply : Branches from IIA , uterine artery , Inf. vesical artery , vaginal artery Lymphatic drainage : Iliac LNs Applied anatomy : - Injury at - Clamping infundibulopelvic ligament - Clamping Uterine artery 1 cm lateral to Cx - Clamping Vaginal angles - Avascular necrosis in meticulous dissection → uretro vaginal fistula , symptoms manifest few days post – operatively	 1) Peritoneum of DP with extra peritoneal fat & cellular tissue 2) Levator ani ms (deep perineal ms) , (urogenital diaphragm) with : - ischiococcygeus - ileococcygeus - pubococcygeus → pubourethralis (decussation around urethra) → pubovaginalis (decussation of ms fibers around vagina) → Puborectalis (decussation of ms fibers around rectum) 3) Perineal ms (superficial) (ischiocavernosus , bulbocavernosus & transverse perineii ms) Fat & skin of vulva <u>* The midline is pierced by urethra , vagina & anal canal</u>	Area of 5 cm between vaginal orifice & anus covered by less hairy skin & less SC tissue Perineal body : - a fibromuscular pyramidal structure between posterior vaginal wall (lower $\frac{1}{3}$) & anterior wall of anal canal  - It is the point of insertion of superficial perineal ms & above it passes the levator ani muscle Applied anatomy : If defective , ie: deficient perineum that causes sexual problem & is treated by posterior colpoperineorrhaphy .



Uterine & cervical ligaments

Support (to prevent prolapse)	Protect important structures
1) Lateral cervical = Mackenrodt's ligament = Cardinal ligament (strongest) Fanning from uterus to lateral pelvic wall 2) Uterosacral (posterior) : From uterus & Cx to periosteum of sacrum 3) Pubocervical (anterior) : From Cx to back of SP	1) Broad ligament: (lateral from uterus to lateral pelvic wall) , Contents : - Fallopian tube (FT) - Parametrial lymphatics & LNs , S & PS nerves - Remnants of Wolffian duct: → Hydatid cyst of Morgagni at fimbrial end of FT → Epoophoron → Paroophoron → Gartner duct : lateral to tube & downward to anterolateral wall of vagina → Paraovarian cyst : in case of cystic dilatation of remnants of Wolffian duct 2) Ovarian ligament (Med between uterus and ovary) : Protects ovarian vessels 3) Infundibulopelvic ligament (Lat between ovary & lateral pelvic wall) : Protects ovarian vessels



NB : Round ligament :

Gubernaculum that attaches cornual end of of uterus to labia majora passing through inguinal canal , it protects Sampson artery & keeps AVF position of uterus.

Sexual differentiation

Gonadal differentiation (Genital ridge)	Internal organ differentiation (Wolffian duct / Mullerian duct)	External organ differentiation
XY XX TDF NO TDF <u>Gonads :</u> = testicles = ovaries <u>Germ cells :</u> Spermatocytes oocytes <u>Sex cord cells :</u> * Sertoli cells W secrete MDIF Granulosa cells Theca cells * Leydig cells → atrophy → Hilar cells	XY Testosterone → growth of <u>Wolffian duct</u> XX NO TDF / NO testosterone * atrophy of Wolffian duct NO Sertoli cells / NO MDIF * growth of <u>Mullerian duct</u> (paramesonephric duct)	XY XX
Anomalies of Ovary	Anomalies of	Anomalies of Hymen
* Aplasia * Hypoplasia * Accessory * Streak gonads (dysgenetic) as in Turner Syndrome	* FT : Aplasia / hypo / accessory osteum / diverticulum * Uterus : Aplasia / hypo / septate / bicornuate / didelphys / unicornuate / arcuate / rudimentary horn * Cx: Aplasia / atresia / incompetent * Vagina : Aplasia / hypo / transverse septum / longitudinal septum	* Imperforate hymen: Failure of canalization of lower part of urogenital sinus

Endoscopy in Gynecology

	Laparoscopy	Hysteroscopy
		
Def	Introduction of an optic lens through <u>umbilicus</u> to visualize peritoneal cavity & pelvic organs	Introduction of an optic lens through <u>cervix</u> to visualize the uterine cavity
Indications	* Diagnostic : → Unexplained infertility → Ch. Pelvic pain / Endometriosis → Cong anomalies of uterus * Operative : → Ovarian (cystectomy / oophorectomy) → Tubal (ectopic / ligation / disconnection) → Uterus (myomectomy / hysterectomy) → Endometriosis : ablation of foci → Adhesiolysis	* Diagnostic: → Infertility → RPL / Ut septum → AUB / Polyp * Operative : → Polypectomy → Septum resection → Myomectomy (submucous) → Division of IU synechia → Tubal occlusion
Technique	- GA - Trendlenberg position (head down) - Veress needle at umbilicus & inflate 3-5 liters CO₂ é pressure 15 mmHg - Introduce lens , light source , camera & manipulator - MB dye may be injected through Cx to visualize patency of FT - Irrigation , evacuation at the end of procedure	- NO anesthesia (in office procedure) , Local or GA in operative procedures - Dorsal lithotomy position - Dilatation of Cx in operative procedures - Uterine distension by CO₂ , glycine (is a must in op procedures) - Lens , light source , camera are introduced - Removal of instruments at the end of procedure
Comp.	- Anesthesia complications / Cutaneous surgical emphysema - Electrosurgical complications to bowel , uterus , nerves - Injury to vessel , intestine , bladder / Infection - Neurological injury in poor patient positioning	- Fluid overload (commonest complication) - Electrolyte imbalance - Anesthesia complications (if used)
Adv.	- Less hospital stay , early return to work - Minimal adhesions - Early recovery , less GIT complications (ileus , gastric dilatation) - Better cosmetic - Rare wound complications (dehiscence & infection)	- Can be done without anesthesia as an office procedure (no dilatation for diagnosis) - Proper visualization of uterine cavity

OBSTETRICS

Bleeding in early pregnancy

1) Abortion **BLEEDING**



Definition : Interruption of pregnancy before viability (< 24 wks) whether fetus is living or dead

Incidence : 15%

Etiology : * Spontaneous abortion → Chromosomal anomalies

* RPL 1st trimester → APS / LPD / ut cavity anomalies / chromosomal anomalies
2nd trimester → Cx incompetence

Types		Threatened منذر	Inevitable حتمي	Incomplete غير مكتمل	Complete مكتمل	Missed منسي	Septic
Clinical picture	S Bleeding	+	++++	++	±	— (Brownish discharge)	Foul vaginal discharge
	S Pain	+	++++	++	—	—	Dull aching pain
	S Abd	= amenorrhea	= amenorrhea	< amenorrhea	Ⓝ sized uterus	< amenorrhea	Tender ut
	S PV(Cx)	<u>Closed</u>	<u>Open</u>	<u>Open</u>	<u>Closed</u>	<u>Closed</u>	Closed or Open
Inv	US	Living fetus	Living or dead	Remnants	Empty ut	Dead fetus (anembryonic sac = blighted ovum)	Usually remnants
	β-HCG	Ⓝ	Not done in emergency	↓↓↓	—	↓↓	↓↓ ++ TLC/ CRP /?ESR
complication		Abortion in 50%	Shock	Septic abortion	±	DIC	Septicemia / septic shock
Ttt		* Rest * PRG supplement	* Correct shock * Termination (SE)	* SE	—	* Termination <u>< 12wks:</u> (no bones) * SE <u>> 12wks:</u> (* bones) medical → PGs If failed medical * hysterotomy	* Broad spectrum antibiotics * Ecboolics Then SE
Give anti-D in Rh-ve ladies							

RPL	
≥ 3 successive spontaneous abortions	Def
* APL (Antiphospholipid \$) * LPD (Luteal phase defect) * anatomical uterine defects (ut septum) (fundal submucous myoma) * ch.anomalies / endocrinal * thrombophilia * Cx incompetence (2 nd Δ) * infections / Trauma	Etiology
* Labs : LA, Endocrinal ,TFTs , anticardiolipin, PRG level * US : eg: Cx length < 2.5cm in Cx incompetence * HSG : eg: funneling in Cx incompetence * Hysteroscopy	Investigations
Ttt of the cause: eg *LD Aspirin LMWH (clexane) SC <u>in APL</u> *PRG in 1 st trimester <u>in LPD</u> *septum resection <u>in septate uterus</u> *myomectomy(hysteroscopic) <u>in fundal submucous myoma</u> *cerclage operation for <u>incompetent Cx</u> (at 12-13wks)	Ttt
 Vaginal common  Abdominal less common Mc Donald / Shirodkar	

2) Ectopic pregnancy

PAIN 😞

Def : pregnancy outside the normal lining of uterine cavity.

Inc : 1.5% (> 95% tubal ectopic)

Et: abnormalities in tube → inflammation / adhesions
→ PRG only contraception / IUD

Pathology :

→ **Tubal :** tubal distension / tubal abortion (tubal mole) / tubal rupture & IPHge

→ **Uterine :** only decidua (absence of villi = Arias stella reaction)

Cl.picture:

- **Symptoms :**
 - **PAIN** , mild vaginal bleeding
 - May present é syncopal attacks / shock
- **Signs:**
 - Gen : ± shock
 - Abd : tenderness at tubal point / “Cullen sign” in case of long standing IPHge
 - PV : Cx motion tenderness (jumping sign = Frog sign) / adnexal fullness/ ut is bulky but < amenorrhea
- **DD :** Acute abdomen , bleeding in early pregnancy

Inv :

- **US :** empty ut & adnexal fullness when β HCG > discrimination zone → TVS = 1500
- **β HCG :** increase but < 60% after 48 hrs (**doubling time**) → TAS = 3500
- ± **laparoscopy :** If done then proceed to ttt via laparoscopy

Ttt:

- **Anti shock measures**
- **Medical ttt:** (MTX 1mg / kg BW **IM**) if :
 - * pt vitally stable / undisturbed ectopic
 - * Ectopic < 3 cm
 - * β HCG < 6000
 - * no fetal pulsations
 - * liver & kidney functions normal
- FU by β HCG at D₄ / D₇ it should ↓ by ≥ 15 %
- **Surgical :**
 - Laparoscopy (if vitally stable) : salpingostomy/salpingectomy
 - Open laparotomy (if vitally unstable) : salpingectomy
- **Anti-D** if pt is Rh-ve

* Other rare types of ectopic :

- in Rudimentary horn
- in ovaries
- in cervix
- over CS scar
- abdominal pregnancy

Ttt: → medical (MTX) : if fulfilling criteria
Or → surgical removal

Inc: 0.15%

3) GTDs (Abnormal proliferation of villi)

Differentiated (With villi)

Undifferentiated (NO villi) (metastatic)

V.Mole (benign)

- Et:** Complete mole : sperm + Anucleated ovum 46 XX
23X — (paternal)
Partial mole : 2sperms + ovum 69XXX/ XXY / XYY
23X /23Y 23X
In extreme Ages , ↓ fat diet , Race (Asia)

Pathology:

Uterus

- Gross: ut enlarged , soft , filled é grape like vesicles é no fetus / sac or liquor (or presence in partial mole)
- Mic: (same in invasive mole too) : differentiated villi

Ovaries: OHSS like (HCG theca lutein cysts)

Clinical picture:

- Symptoms:** vaginal bleeding / prune juice discharge
lower abdominal pain / enlargement
- Signs:** Gen : anemia / thyrotoxic (TSH like) /
↑BP (HCG effect)
Abd : Ut > period of amenorrhea /soft Doughy ut
PV& bimanual : soft Cx / vesicles / full adnexa
- Complications:** Invasive mole 20% / Chorioca. 5%
- DD:** bleeding in early pregnancy

Inv:

To confirm diagnosis : US: snow storm appearance /
Ov theca lutein cysts

- For follow up : β- HCG : very high level
- To detect complications : CXR / MRI pelvis

Ttt:

Suction Evacuation (é ecbolics & antibiotics)
FU é β- HCG weekly till -ve then monthly for 1 year
(use COCPs as contraception)

NB:

- In old age , no desire for fertility = TAH
- If plateau or rising HCG : use MTX
- Anti - D in partial mole in Rh -ve non sensitized pts

GTT (GTN)

Invasive Mole (locally m)	Chorio Ca.	Placental site T.	Epithelioid T.
Low risk		High risk	
* < 4 months from preg. * No liver or brain M. * No prev chemo ttt: Single agent chemotherapy (MTX) good prognosis		* > 4 months from preg. * Liver or brain M. * Previous chemo ttt: Combined chemotherapy (MAC) bad prognosis	

Pathology:

- Gross: enlarged ut é friable tissues é areas of hge & necrosis
- Mic: undifferentiated villi
- Grading :
 - G1: < 5% malignant undiff.cells
 - G2: 5-50% malignant undiff.cells
 - G3: > 50% malignant undiff.cells
- Spread:
 - Direct : to surrounding
 - Lymphatic : very rare
 - Blood : Lung B L B

Cl.pict

- ⑤: vaginal bleeding é H/O of pregnancy event
- ⑤: Gen : anemia / metastasis
- Abdominal : enlarged soft uterus / metastasis
- PV& bimanual : enlarged ut / adnexal fullness / vag.masses
- FIGO staging
 - Stage I : confined to uterus
 - Stage II : local spread (ut / ov / vagina)
 - Stage III : + lungs
 - Stage IV : distant other than lung

Inv:

- To confirm diagnosis : HCG / US
- To detect spread : Chest X-ray / MRI pelvis
- To assess fitness of pt for ttt: LFTs / KFTs

Ttt:

- Low risk: single agent chemo (MTX) é FU 1year after -ve HCG
- High risk: combined chemo (MAC) é FU 1year after -ve HCG

NB: Hysterectomy if:

- * Cases complicated by severe bleeding
- * Chemoresistant cases as placental site & Epithelioid tumor


4) Associated Gynecologic conditions

Bleeding in late pregnancy (APHge)

Definition : Bleeding after fetal viability (24 wks) till delivery

Etiology :

- Maternal**
 - Obstetrics : Abruptio-placentae (accidental hge) / Placenta previa (PL PRV)
 - Gynecology : Vaginal lesion / Cervical lesion
 - Systemic : Drugs (anticoagulants) / Systemic disorders
- Fetal :** vasa previa (minimal bleeding → severe fetal distress)

	Accidental Hge (commonest)	Placenta previa (2 nd most common)
Definition	 Normally implanted placenta	 Placenta implanted on LUS
Risk Factors	PG (PIH) / External trauma / APS /Smoking Sudden ROM in polyhydramnios	Multipara / old age / Prev CS / prev PL PRV Malpresentations / Twins
Mechanism (Pathology)	* In PIH: Blood → inbetween myometrial fibers → tender hard ut → Couvelaire ut (concealed accidental Hge) ▶ Atonic PPHge * In trauma: soft ut & normal myometrium (revealed accidental Hge)	Shearing mechanism between LUS & placenta → <u>Unavoidable</u> bleeding
Types	Mixed / Revealed / Concealed	Complete centralis / incomplete centralis Marginalis < 2 cm from os / Lateralis > 2 cm from os
Cl.picture	Symptoms	±Vag bleeding usually abdominal pain
	General	PIH ? ± Shock (BP may not be ↓)
	Abd	Tender ut (in concealed) / FL > amenorrhea
Signs	PV & speculum	No ut tenderness / FL = amenorrhea / malpresentations
	Contraindicated till exclusion of PL PRV	
Complications	HELLP / DIC / Atony / Sheehan syndrome	Placenta Accreta / Increta / Percreta (PAS)
Investigations	Maternal : US /CBC / LFTs / KFTs / Inv for DIC / HELLP Fetal : Assessment of fetal wellbeing (DFMC / NST / BPP / Doppler)	

Management of APHge:

Termination if :

- * Reached maturity
- OR * Labor pains
- OR * Maternal or fetal complication

Otherwise **Conservative** until one of 3 factors above are reached

Termination

VD
In Accidental hge
(unless CI)

LSCS
In PL PRV
(unless Lateralis)

* In case of abnormal adhesions of placenta, with no plane of cleavage (Increta / Percreta) → **ttt** : Cesarean hysterectomy (= USCS followed by hysterectomy) as any attempt to separate the placenta increta / percreta will result in severe bleeding

NB: *Resuscitation in shock (if present)

* **ttt of complications if present ie DIC / HELLP / Atony**

NB: Placental migration : < 24 weeks

Lower edge of placenta may be seen near int os , but é formation of LUS , placenta migrates upward to its normal position, away from internal os & the condition resolves in > 90% of cases

Intrapartum Hge

= **Obstetric Trauma**

Rupture uterus

Risk F : MG / scarred ut / oxytocin , PG /Instruments

Cl picture (S) → Pain relieve
(S) → Abd : Fetal parts easily felt
FHS inaudible
→ PV : head reced upward

Management :

* Resuscitation
* Laparotomy → Repair or
→ Subtotal hysterectomy

	Cervical tears	Perineal tears
Due to	Improper management of 1 st stage	Improper management of 2 nd stage
Risk F	Echbolics /instruments / PPT labor / manual dilatation/ macrosomia	Omitting episiotomy / malpresentations/ Macrosomia
Types	Lat / ant lip	1 st : skin 2 nd : skin + superficial perineal ms 3 rd : 2 nd + anal sphincter 4 th : 3 rd + rectal mucosa
ttt	Prevention by proper management of labor Repair ± replacement	Repaired in layers + post operative care

Old complete perineal tear

Def : Missed & unrepaired 3rd degree tear at delivery

CP : (S) Incontinent to flatus / stool

(S) → defect in perineal body
→ Red rectal mucosa may be seen
→ 2 dimples of fibrosed retracted anal sphincter
→ Absent corrugation of anus sp posteriorly
→ PR exam : no tone of levator

Ttt :

* Prevention by early diagnosis & ttt at time of labor
* Preoperative preparation
* OP : H shape incision / repair in layers / + post colpoperineorrhaphy
* Post Op care *ttt of cause

Complications of 3rd stage

I) PPHge (bleeding after delivery) 🩸

1ry: 1st 24hrs

2ry: > 24hrs till end of puerperium due to infection /retained parts / choriocarcinoma /other Gyn causes

1ry PPHge

Def : bleeding > 500cc in VD / > 1000cc in CS

Inc : leading cause of maternal mortality in Egypt

Et :

- * Atony (overdistension , prolonged labor , ecbolics, anemia ,...)
(most common)
- * Traumatic(Cx , perineal , vag laceration due to instrumental or macrosomia)
(2nd common)
- * Retained placenta (remnants) (3rd common)
- * DIC or systemic disorders
- * Ut inversion (may not present é bleeding but pain)

Cl.picture :

- S : → Gen : anemia / shock
→ Abd : lax ut (atony) / firm (trauma) / rising (bleeding)/
inverted cup (inversion)
→ PV : dark blood clots (atony) / fresh bleeding (trauma) / lacerations /
retained placenta / inverted ut

Complications : Death / Sheehan \$...

Inv : CBC (inv for complications)

Ttt:

A) PREVENTION

(proper ANC / 1st , 2nd , 3rd stage management)

B) Actual 1) Ecbolics + massage + bimanual compression

Vag { 2) Exploration of genital tract for trauma
3) Intrauterine Bakry ballon (placed vag)

Abd { 4) Bilat uterine / IIA ligation
Vs
4) B-Lynch suture
5) Subtotal hysterectomy

NB : Resussitation while doing these measures

II) Retained placenta

Def : placenta didn't deliver within 30 min of fetal delivery

Et : → Retained separated (atony / rupture ut / constriction ring (rare))
→ Retained non separated (PAS)

Ttt: 1) Ecbolics + Brandt - Andrew's 2) Manual separation
3) If only parts = D&C by ovum forceps 4) ttt of cause (if present)

NB: in PAS

* **Accreta (80%) :** there is usually a plane of cleavage so separation of placenta occurs (take care if increta / percreta)

* **Increta / Percreta :** planned cesarean hysterectomy ie classical CS, removal of fetus , closure of wound & proceeding to TAH with ovarian preservation éout any attempt to separate placenta (to avoid severe bleeding)

III) Uterine inversion

Def :

Et : traction on placenta while uterus is lax

Cl.picture : *Gen : neurogenic shock
*Abd : absent fundus
*PV : inverted cup (decidua seen)

Ttt: *Avoid

* **Actual :** Manual reposition **or** Hydrostatic pressure reduction

IV) Shock (Hgic / Neurogenic / Septic)

Hgic shock

Def : state of circulatory impairment

Cl.picture :

- 1) Amount of blood loss (< 15% , 15-30 % , 30-40% , > 40 %)
- 2) HR (< 100 , 100-120 , 120-140 , > 140)
- 3) BP (N or ↓)
- 4) Urine output (> 30ml/ hr , 15-30 , 5-15 , < 5ml/hr)

Ttt: Airway (maintained patent)
Breathing (oxygenation)
Circulation (adrenaline / perfusion)

Replacement :

- * Whole blood
- * Fresh Frosen Plasma (all clotting factors)
- * Cryoprecipitate (factors VIII , XIII , Fibrinogen)
- * Packed RBCs
- * Platelets

Septic shock

Ttt : Broad spectrum antibiotics + ttt of caus

DIC

Def : consumptive coagulopathy / hypofibrinogenemia

Et :

- * Massive blood loss * Massive transfusion * Concealed accidental Hge
 - * HELLP * Amniotic fluid embolism
 - * IUFD → ++ **extrinsic pathway** (thromboplastin release)
- } ++ **intrinsic pathway** (endothelial cell damage)

Cl.picture : of cause / bleeding from venipuncture

Inv :

- ↓ fibrinogen < 100 mg / dl
- ++ FDPs (++ D-dimer)
- ++ PT , PTT

Ttt :

- Of cause
- Replacement (FFP / Cryoppt)
- Never ×× Heparin

Amniotic Fluid Embolism (AFE)

Def : triad of hemodynamic , respiratory compromise & DIC (no fever)

Risk F : PPT labor / induction & augmentation / pl abruption

Ttt : CPR / intubation / inotropes / FFP / avoid excess fluid

Maternal collapse

Def : acute cardio respiratory , CNS failure → Cerebral hypoperfusion → death

Et : hemorrhage / thromboembolism / AFE / ICHge / cardiac disease / sepsis / anaphylaxis

Ttt : immediate CPR, oxygenation & circulatory support

Maternal mortality

Et : death by obstetric triad (preventable)

- Hemorrhage
- Preeclampsia
- Infection

Ttt: prevention by proper ANC & IPC


- Other causes :
- Thromboembolism
 - Cardiopulmonary
 - AFE

Female bony Pelvis

Anatomy of Fetal skull

Terminologies

- Inlet




- Cavity

Plane of least pelvic dimensions

Plane of greatest pelvic dimensions
12.5 cm


- Outlet



Outlet segment

Engaging diameters

A / P engaging diameters:



Transverse engaging diameters:

- BPD (largest) = 9.5 cm
- Supra parietal / Sub parietal = 9 cm
- (in asynclitism)
- BTB = 8 cm
- BMD = 7.5 cm


Engagement: passage of widest transverse diameter of the presenting part (eg :BPD) through plane of pelvic inlet.

(Rule of 5: 2/5 = engagement)

Lie: (longitudinal / transverse)

Attitude: fetal parts (flexed / extended)

Presentation: 1st felt on PV



Denominator :

Bony landmark of the presenting part

Asynclitism (tilt) : (eg : ant asynclitism = post parietal bone presentation , when sagittal suture is toward ant.)

Position:

Relation of fetal denominator to the right / left & ant /post of the mother

Station:

0 when occiput is at the level of ischial spines in vertex presentation = Engagement

Effacement : LUS is taken in UUS

Leopold manoeuvre:

(F.level / F.grip / UG / 1st & 2nd PG)

Naegle's formula :

LMP + 7days + 9months → EDD

Obstetric code:

* 4 digit code

F	P	A	L
> 36 wks	24-36	< 24 wks	now

* GPL

* P +

Normal labor

Def : passage of single / living / cephalic / vertex / term / through birth canal / éout interference / é no complications (maternal & fetal) / éin 4 – 24 hrs

Et (Theories): ++PGs / ++ fetal cortisol / ++ ut distension / -- PRG / -- pl oxytocinase

Stages	1 st stage Cervical changes (effacement & dilatation) Latent < 4 cm Active > 4 cm	2 nd stage (Fetal delivery)	3 rd stage (Placental delivery)
Duration	12 -18hrs in PG 6 - 8hrs in MG	2hrs in PG 1hr in MG	Up to 30 min in both
Mechanism	Ut contractions (true labor pains) → Cx effacement & dilatation	- Descent - Engagement - ↑ Flexion - Int rotation - Extension - Restitution - Ext. rotation - Expulsion may occur in late pregnancy	- Placental separation, detected through : * Gush of blood * Elongation of cord * Suprapubic bulge - Placental descent - Placental expulsion: * Schultze * Duncan
Aim	To get <u>strong efficient</u> (3 cont/ 10 min / each lasting 1 min / é intensity 40-60 mmHg) to cause dilatation of 1.2 cm/hr in PG Or 1.5 cm/hr in MG	To allow extension <u>after</u> crowning to distend vulva by <u>SOE</u> (↓↓ perineal tears)	To minimize bleeding (500cc)
Through	↑↑ contractions by : AROM / oxytocin ↓↓ contractions by : stop oxytocin / rehydration / epidural	Ritgen maneuver (perineal support)	do <u>active</u> management through use of <u>ecbolics</u> (eg ergometrine/oxytocin)
Management	- H/O: medical / surgical / obstetrics F P A L LMP, EDD, GA - Exam : → Gen : vital signs → Abd : Leopold maneuvers / FHS → PV : presentation / position / Cx dilatation / effacement / station / ROM - Nutrition ? IV fluids / Analgesics / Enema - Partogram : for FU progress of labor	- Lithotomy / sterilization / towling - Instruct her to bear down during contractions& relax inbetween - Maintain flexed occiput & support perineum	- Wait for signs of separation - Brandt – Andrews maneuver - Explore placenta & memb after their delivery for completion - Explore birth canal for Tears
NB:	External <u>CTG</u> ONLY in high risk pregnancy (augmentation is a high risk)	<u>Episiotomy</u> ONLY if indicated	- 2hrs later observe for complications (4th stage) - Give Anti-D in Rh –ve

Malposition & Malpresentations

	Occipito - posterior	Face
Def	Longitudinal lie in which head is flexed & occiput posterior (occiput is denominator) {It's a malposition & not a malpresentation}	Longitudinal lie in which head is fully extended (mentum is denominator)
Inc	20% at time of labor	1/500
Position	ROP : most common occupying wider Right oblique diameter	LMA : most common as it results from extension of ROP
Et	Gen: CPD / Fetal anomalies / MFG / ... Specific : Android / Anthropoid pelvis	Gen : CPD / Pendulous abdomen / ... Specific : Anencephaly
Mechanism	* correction of deflexion 90% → long ant rotation → DOA & delivery by extension as OA * mild deflexion 1% $\xrightarrow{\text{occiput 1st}}$ ant Rot $\frac{1}{8}$: (deep transverse arrest) obstructed * moderate deflexion 3% $\xrightarrow{\text{both occiput \& sinc}}$ NO rotation : (persistent oblique OP) obstructed * severe deflexion 6% $\xrightarrow{\text{sinc 1st}}$ post Rot $\frac{1}{8}$: DOP (face to pubis) & delivery by flexion	* MA : ant rotation → DMA & delivery by flexion * MP $\begin{cases} \rightarrow \frac{2}{3} \text{ long ant rotation to MA \& delivery by } \text{flexion} \\ \rightarrow \frac{1}{3} \text{ no rotation or post rotation to DMP : } \text{obstructed} \end{cases}$
Comp.	Maternal : Prolonged labor / PROM / PPHge / Puerperal sepsis Fetal : Distress / Asphyxia / Birth injuries / Instrumental delivery / Comp of associated anomalies	
Management	Pregnancy : Leopold : as OA : FL (same) / FG (buttocks) / UG (back Rt) / 1 st Pelvic Grip (done : delayed engagement) / 2 nd Pelvic Grip (not done) Auscultation : FHS below umbilicus / US	
	Labor: PV : occiput 1 st stage : watchful expectancy for <i>factors that favor long anterior rotation</i> : (Roomy pelvis (no CPD) / Good pelvic floor muscles / Strong uterine contractions / Adequate liquor) *90%: as OA (assess factors for long ant rotation) *6% : Face to pubis (distending diameter OF 11.5 cm) * do episiotomy *4% $\begin{cases} \rightarrow \text{Kielland for rotation \& extraction} \\ \rightarrow \text{Ventouse for rotation \& extraction} \end{cases}$ } If no CPD or Safer CS	PV : chin (Tumefaction) *MA : delayed engagement + episiotomy (SMV 11.5 cm) *MP $\begin{cases} \rightarrow \frac{2}{3} \text{ factors for long ant rotation (as MA)} \\ \rightarrow \frac{1}{3} \begin{cases} \rightarrow \text{Forceps delivery (not done)} \\ \rightarrow \text{NEVER ventouse (CI)} \end{cases} \end{cases}$ or Safer CS
Brow		
Def: Head is midway between flexion & extension Inc : 1/2000 Et : general causes		
Mechanism : $\begin{cases} \rightarrow \text{Transient brow :delivery as OP or MA} \\ \rightarrow \text{Persistent brow (MV = 13.5 cm) : } \text{Obstructed} \end{cases}$		
Management : $\begin{cases} \rightarrow \text{Transient brow : manage accordingly} \\ \rightarrow \text{Persistent brow : CS} \end{cases}$		

Breech presentation

Def : It is a longitudinal lie in which buttocks with feet (complete) / buttocks only (frank) / feet (footling) / knee are the presenting part.

Incidence: 3.5% at full term / 25% at 28 wks

Etiology: → **General :** contracted pelvis / ut septum / fibroid / pl.previa / MFG
→ **Specific:** Hydrocephalus (in full term) , Prematurity

Position: LSA

Types: → Complete (MG)
→ Frank (PG)
→ Footling / knee ...

Mechanism :

for → Buttocks (Descent / Eng (BTD 9.5cm) / Int rotation / ant buttock hinge below SP, post buttock deliver 1st by Lat Flexion of spine)
→ Shoulders (Descent / Eng (BAD 12 cm) / Int rotation / post shoulder deliver 1st by Lat Flexion of spine)
→ After coming head (Descent /Eng (BPD 9.5) / Int rotation(opposite direction as it enters pelvis in opposite axis)/ delivery by Flexion)

Complications:

→ **Maternal : 4 Ps** (Prolonged labor / PROM / PPHge / Puerperal sepsis)

→ **Fetal** → Retained after coming head
→ Post. rotation of the head →do→ Prague manoeuvre
→ Extension of arm →do→ Löwset manoeuvre
→ Fetal birth injuries
(nerves: Erb's palsy / bones : hip dislocation / viscera: rupture spleen,rupture liver, rupture anal sphincter& hymen defloration)
→ Fetal distress / Sudden compression & decompression

Diagnosis : → **During pregnancy** → Leopold maneuver (FL:same as amenorrhea period / FG: head / UG / PG: buttocks)
→ FHS (above umbilicus)
→ US
→ **During labor** + PV (tip of sacrum + ischial tuberosities at same plane)

Management:

→ **During pregnancy:** ECV (60% success) , at 36 wks if not CI

Side effects: 1) ROM /cord prolapse 2) Pl. separation
3) Loops of cord around fetal neck
4) Fetal distress 5) ++ labor

→ **During labor :** (Assess for unpredictable hazards sp in PG)

→ VD: *Assisted breech delivery : spontaneous delivery of buttocks & shoulders , BUT Assisted delivery of head →
Burns-Marshall Mauriceau-Smellie-Veit ± Kristiller's manoeuvre Piper's forceps
(Leave baby hanging by its weight) (Jaw flexion shoulder traction) (supra pubic pressure on head by assistant)

*Breech extraction **ONLY** done in : 2nd twin if breech / fetal distress é fully dilated Cx and NO CPD

→ CS

Indications of CS: 1) Extended neck / 2) Twins 1st breech (locked twins) / 3) Preterm / 4) PG (80%) ??

5) weight < 2.5 kg , > 3.5 kg / 6) Breech presentation other than complete or frank / 7) Other indications for CS

Shoulder presentation

Def : Transverse Lie é scapula as denominator (ant or post)

Incidence: 0.5%

Etiology: ➔ **General :** CPD / Septate ut / Pl. previa
➔ **Specific** Pendulous abdomen in multipara

Position: ➔ LSA
➔ RSA

Mechanism : NO mechanism (Obstructed labor)

Complications: ➔ **Maternal** → 4 Ps
➔ **Fetal** + neglected shoulder

Diagnosis : ➔ **During pregnancy** ➔ Leopold manoeuver (FL → less than period of amenorrhea / FG → empty)
UG → head on one side & buttocks on the other side / 1st PG → empty)
➔ FHS → at level of umbilicus at the head side
➔ US

➔ **During labor:** + **PV** → Grid – iron (ribs)

Management: ➔ **During pregnancy:** ECV till 1st stage of labor
➔ **During labor :** **CS**

NB : IPV and breech extraction is ONLY done for 2nd twin (if transverse)

NB : neglected shoulder : Arm prolapse with ROM for long time & IUFD , delivery by **CS** (mostly upper segment)
for maternal safety to prevent rupture uterus

é intact memb.

← Cord presentation & Cord prolapse →

é ROM

Obstetric emergency

Def : cord is below presenting part either é intact membranes (presentation) or é rupture membranes (prolapse)

Et : general : ➔ **Maternal**
➔ **Fetal** (specific) : malpresentations as shoulder / incomplete breech / footling / ...

Diagnosis : **PV** : mind PULSATING or NOT

Management : ➔ **IF PULSATING :** **urgent CS** , while transferring her to OR: reposit the cord , give O₂ , knee chest position
or trendelenburg position / **IF fully dilated Cx:** ➔ forceps in cephalic engaged
➔ breech extraction in breech presentation
➔ **IF NON PULSATING :** allow VD (not an emergency)

Complex presentation

Def : **Arm** beside head or buttocks in 2nd stage

Management : reposit the arm & proceed é VD

Unstable Lie

Def : Fetus continuously changes its position & / or presentation after 34 weeks GA

Management : stabilized induction ie induction of labor while it is in a cephalic presentation after 36 wks

Multifetal gestation (Twins in 97% of cases , Triplets in 1-2%)

Definition: Simultaneous presence of 2 or more fetuses in the uterus

Inc: DZT 1-2% > MZT $1/250$

ET: → MZT (1 sperm + 1 ovum)

cleavage in MZT

- If < 3 days → Dichorionic Diamniotic 30%
- If 4-8 days (chorion formed) → Monochorionic Diamniotic 65%
- If 8-12 days (ch & amnion formed) → Monoch. Monoam. 5%
- If > 12 days (parts formed) → Conjoined (rare)

MOST SERIOUS



→ DZT(2 sperms + 2 ova) é induction of ovulation

Cl. picture : → **S :** large sized abdomen / ++ S of pregnancy

- **S :**
 - General : ++ anemia / PIH / HEG / GD / ...
 - Abdominal : Leopold maneuvers (FL / FG / UG / 1st Pelvic)
 - PV: presenting part small in comparison to large abdomen

FHS : galloping sign

Inv: US

Mechanism : Depends on the presentation of 1st & 2nd fetuses

Complications:

→ **Maternal :** 4 Ps + ↑↑ rate of medical disorders (hyperemesis / HTN / anemia / GD / ...) + pressure symptoms

→ **Fetal** 1) vanishing twins / fetus compressus / fetus papyraceous

- 2) MZT → MC , MA & ttt : discordant twins (shared placenta)
 - One fetus LGA / polyhydraminos / polycythemia
 - Other fetus SGA / oligohydraminos / anemic
- ttt : laser ablation of anastomotic vessels
- Conjoined twins (due to delayed cleavage)

3) locked twins , if we allow VD when 1st fetus is breech (which should not be done)

4) death of one fetus : 1st trimester (early) : Little risk / 3rd trimester (late) : ++ DIC

5) PTL (COMMONEST complication) & prematurity complications

Management : → **During pregnancy :** more frequent ANC visits due to more ++ of medical disorders during pregnancy

- **During labor :**
 - CS : if 1) other indications of CS 2) > 2 fetuses
 - 3) monoamniotic twins or conjoined twins 4) 1st fetus is non cephalic
 - VD : if 1st fetus is cephalic & no other indications of CS

Delivery of 2nd twin

- If cephalic (most common)
 - ROM &
 - Allow VD spontaneous
- if breech
 - ROM &
 - Do breech extraction
- if transverse
 - ROM &
 - Do IPV & breech extraction

Guard against Atonic postpartum hge by prophylactic ecbolics

OVD (operative Vaginal delivery)

	 Forceps	 Ventouse (vacuum extractor)
Start	By Chamberlen family (1 st to use forceps on <u>living</u> fetus) in year 1560-1730	By Malmstrom in 1954
Benefits / Characteristics	* In modern obstetric , only <u>low</u> (outlet) forceps is used <u>for extraction only</u> * Kielland long forceps is rarely used for rotation & extraction in OP * in fetal distress to shorten 2nd stage * In Face presentation & PTL * Aftercoming head of Breech (Piper's forceps) * Dead fetus	* For rotation & extraction in OA or OP positions * Less encroachment on maternal pelvic space * Time consuming (need 20minutes to induce –ve pressure) * not used in fetal distress * CI in face or preterm (only used in <u>vertex</u> presentation) * Only used on <u>living</u> fetus as –ve suction & scalp edema formation ie chignon (which is the mechanism of action of ventouse) needs living fetus to be formed
Pre-requisites	Fully dilated Cx / Engaged head (ie station 0 at least) / NO CPD / Membranes ruptured / Presence of uterine contractions / Empty bladder & rectum / Antiseptic techniques / Anesthesia	
Complications	Maternal (mainly) : - lacerations & tears (perineal / vaginal / cervical & even rupture uterus) - PPHge (traumatic or atonic) Fetal : - if wrong application : head compression , skull fracture & ICHge - cephalhematoma (bone fracture or fissure) - facial nerve injury	Fetal (mainly) : - cephalhematoma - scalp lacerations - ICHge if excessive –ve pressure in preterm fetus with fragile BVs Maternal : - If wrong application (includes cervical tissue in the ventouse cup) → Lacerations

Cesarean Section

Def : Delivery of a viable fetus through an abdominal & ut incisions.

Types of CS :
 ↳ **LSCS** → transverse incision (by default)
 ↳ Vertical incision (in certain situations)
 ↳ **USCS (Classical CS)** usually in PAS

Indications:

↳ **Maternal :** CPD / previous CS
 ↳ **Fetal :** Malpresentations / fetal distress

Techniques:

	LSCS	USCS
Uses	Usually done	In placenta accreta
Inc of rupture	0.2 – 0.9 %	×10 times (2-9 %)
Scar	Formed of 2 ms layers ↳ less Hge / better coaptation / less hematoma ↳ better healing (strong scar)	Formed of 3 ms layers ↳ more Hge / less coaptation / more hematoma ↳ worse in healing (weaker scar)
Peritoneum	Better peritonization & less adhesion formation	Visceral peritoneum is adherent ↳ more adhesions later

Complications of CS :

↳ **Intraoperative :**
 * Anesthesia complications
 * Injury to bladder/ intestine / BVs
 ↳ **Early post operative:**
 * Reactionary Hge
 * Ureteric injury symptoms / fistula
 ↳ **Late post operative :**
 * Wound infection
 * Paralytic ileus , acute gastric dilatation
 * Thromboembolic complications
 * Adhesions & subsequent tubal & peritoneal factors of infertility
*** Placenta accreta if implanted on the scar site**

NB : VBAC (Vaginal Birth After CS) or TOLAC (Trial Of Labor After CS)

Prerequisites :

- Only one previous LSCS
- Never after USCS (classical CS), hysterotomy or myomectomy scars
- No current indication for CS

Complications of VBAC : rupture uterus

“ Previous normal VD followed by CS improves the chance of a safe and successful VBAC ”

Episiotomy

Definition : an incision in post vaginal wall , perineum & skin done during vaginal delivery to widen the vulval introitus for the fetus

Indications:

Maternal :
 ↳ rigid perineum
 ↳ instrumental delivery

Fetal :
 ↳ macrosomia
 ↳ malpresentations or malposition
 ↳ prematurity (to avoid sudden compression & decompression of fetal head)

Timing : Just before crowning

Types :

	Median	Mediolateral
Benefit	More anatomical More cosmetic Less dyspareunia Less pain Less blood loss Easier repair Better healing	No extention to anal sphincter & rectum so avoid damage of anal sphincter (ONLY advantage over median episiotomy)

Complications :

- extension to anal sphincter in median type
- hematoma → Infection
- later dyspareunia

Analgesia & Anesthesia

Anesthesia in labor

Spinal is anesthesia of choice for CS : in subarachnoid space (hypotension & headache , ↳ IV preload is needed)

Epidural :

(hypotension , ↳ IV preload is needed)

General : IV anesthetics as thiopental Na (ketalar) , in addition to inhalation gas drugs as N₂O & O₂ (affect fetus & mother)

Analgesia in labor

I) Pharmacological

General	Local
1) Narcotics: *Pethidine × 10 *Morphine × 5 *Butorphanol Side effect: Neonatal RDS if given < 2hrs before delivery Antidote : naloxone (Narcan)	1) Epidural : used all through labor, ie: intrapartum & postpartum Side effects : * block motor too (not just sensory) * loss of urge of straining * accidental puncture of dura : headache
2) Non- Narcotics : * Benzodiazepines: diazepam (valium) * Phenothiazine derivatives	2) Local infiltration anesthesia (é lidocane) Most commonly used (for episiotomy or tears)
3) Inhalation : * N ₂ O + O ₂ → 50:50 (laughing gas) * Trilene (obsolete due to its toxic metabolites)	3) Pudendal nerve block At level of ischial spine , injected through the vagina
	4) Paracervical block * Injected on either side of the cervix through the lateral fornices * Side effect : fetal bradycardia (↳ rarely used)

II) Non-Pharmacological

- 1) ANC classes (how to relax / breathing excercises / abdominal & pelvic floor ms exercises)
- 2) TENS (Gate theory of pain)
- 3) Acupuncture (Gate theory of pain)
- 4) Water birth

Fetal surveillance (assessment of fetal wellbeing)




Antepartum

(from viability till labor)

1) **DFMC** : (N) 10 movements / 12 hrs

2) **NST** : fetal heart acceleration in relation to its movements (for 20 min)

(N) 15 b/min acceleration for 15 sec (= reactive NST)



- If non reassuring NST → Repeat for another 20 min
- If non reactive NST → * Do BPP

3) **BPP** : US to detect (movement / tone / breathing / AFI)

Each item → 0 if not present
→ 2 if present

* Total score = 8

- If < 6 / 8 * do doppler

(NB: Modified BPP = NST + AFI)

4) **Doppler study**: for umbilical artery , MCA

- If ++ resistance (low flow) * placental insufficiency
- If NO flow * Alert sign
- If reversed flow * Action should be taken

Consider termination: IOL OR CS

NB : If high risk cases * assessment of fetal wellbeing

DFMC → done daily

NST / BPP → done biweekly

Doppler → done weekly

Intrapartum

(in 1st & 2nd stages of labor)

1) **Passage of Meconium (in cephalic presentation)** :

indicates fetal distress

2) **CTG** : relation of FHR to uterine contractions

- (N) FHR = 110 – 160 b / min

- If < 100 bradycardia
 - If > 160 tachycardia
- } = fetal distress

CTG ONLY FOR HIGH RISK

- Loss of beat to beat variability = fetal acidosis

- Early deceleration = head compression (N)

- Variable deceleration = cord compression (exclude cord prolapse)

- Late deceleration = fetal distress (sp if persistent late deceleration)

3) **Fetal scalp PH** (N) PH = 7.25 – 7.35

* If 7.2 – 7.25 → * mild acidosis

* If < 7.2 → * severe acidosis

4) **Scalp stimulation** : → FHS acceleration

5) **Vibroacoustic stimulation** : → FHS acceleration

6) **Internal (direct) electronic monitoring** :

To record fetal P wave , QRS complex & T wave

Management of Intrapartum distress (asphyxia)

* Stop oxytocin / give IV fluids

* mask O₂ / put pt in Lt lateral position

Persistent distress

Immediate Termination

Returns back to (N)

continue labor & close observation

Not fully dilated

CS

fully dilated

VD

→ Forceps in cephalic pr.
→ Breech extraction in breech pr.

Abnormal labor

Def : labor lasting < 4hrs or > 24 hrs due to abnormalities in passages (pelvis) / passenger (fetus) / power(ut. contractions)

Types:

1) Precipitate labor

Def : < 4 hrs for 1st, 2nd & 3rd stages

Et : multipara / small fetus / roomy pelvis / strong contractions

Comp: * Maternal: lacerations of Cx / P.sepsis / PPHge

* Fetal : fetal birth injuries

Ttt: → If seen before delivery : epidural to slow down progress / Perineal support in 2nd stage

→ Explore birth canal for lacerations / antibiotics prophylaxis /neonatal exam for birth injury

2) Prolonged labor

Def : delay in progress <1cm /hr in PG or < 1.5cm /hr in MG

“While arrest of labor : no progress at all (in Cx changes or fetal descent) for **1 hr in MG** or **2 hrs in PG** in the presence of strong efficient ut contractions whether in active 1st or 2nd stages”

NB: Double this time if epidural is used

Et : → CPD

→ Hypo / Hypertonic ut. Inertia

Comp: * Maternal: exhaustion / PPHge / P.sepsis

* Fetal : distress / asphyxia

Ttt : → If CPD : CS

→ If hypotonic : amniotomy / oxytocin augmentation

→ If hypertonic : stop oxytocin / IV fluids / epidural

NB: In 2nd stage arrest , assess whether it is due to exhaustion (≠ use instrumental delivery if engaged) or obstruction (≠ do CS)

3) Obstructed labor

Def : arrest of 2nd stage (although presence of strong uterine contractions) due to CPD.

Cl. picture : * Gen : dehydration / exhaustion

* Abd : pathological retraction ring (Bandl's ring) DD from contraction ring

Pathological retraction ring (Bandl's ring)	Constriction / contraction ring
* Between UUS & LUS	* Spasm of circular smooth ms fibers
* Moves upward	* Any site
* Seen abdominal & felt vag	* Felt vag only
* + Fetal distress / severe maternal exhaustion	* No maternal or fetal distress
* Relieved by CS	* Ms relaxation / anesthesia

* PV : caput / dry vagina / oedematous / fully dilated Cx / station -2 -1

Comp : * Maternal : rupture ut / necrotic VVF / PPHge / P. sepsis

* Fetal : distress / asphyxia

Ttt: delivery by CS (NO instrumental as head is NOT engaged)

NB : PG responds to obstruction by atony while MG responds to obstruction by rupture uterus

4) Cx dystocia

Def : rigid Cx that doesn't dilate or efface in the presence of efficient ut .contractions

Et : spasm / scarred Cx

Ttt: spasmolytic / CS (to avoid annular detachment of Cx or rupture ut)

5) Shoulder dystocia

Def : Failure of delivery of shoulder after head delivery due to impaction of anterior shoulder at the symphysis pubis

Et : fetal macrosomia

Cl.picture: Turtle sign

ttt: → Prevention by CS when indicated

→ Actual :

Shoulder dystocia drill: (call for help / generous episiotomy / Mc Roberts (hyperflexion of maternal thigh) ± suprapubic pressure)

Other measures that may be tried if above faild:

* Wood Cork Screw * Post. Shoulder delivery * Rubin maneuver

Other measures :

* Symphysiotomy * Zavenilli Maneuver * # ant clavicle * post axilla sling * Cleidotomy: # clavicle in a dead fetus

CPD

Def : one or more of the pelvic diameters is decreased so as to interfere é normal vag delivery

Et : diseases or trauma to spine , femur or pelvic bones of the mother

Cl.picture :

* H/O

* Exam → Gen: for gait / height
→ Abd : pelvimetry
→ PV : Internal pelvimetry & CPD tests:
1) Pinard : NO PV
2) Muller-Kerr : é PV

Ttt: → If no CPD : allow VD
→ Moderate CPD (1st degree) : trial of labor
for undetermined factors (moulding of head / yielding of pelvis / strong ut cont.)
→ Severe CPD (2nd degree): CS

PROM

Definition : ROM after fetal viability & before onset of labor

Incidence : 10%

Etiology:

- Idiopathic
- Infection (commonest) : GTI , UTI
- Cx incompetence
- Polyhydramnios
- Local membrane defect / smoking



Complications :

Maternal : infection : Chorioamnionitis (most serious)

Fetal: Preterm labor within 24-48 hrs (commonest)

Clinical picture :

Symptoms : gush of fluid

Signs :

- General : fever (in case of infection)
- Abdominal : FL < period of amenorrhea / tender uterus in chorioamnionitis
- PV(under **STRICT** aseptic conditions) : speculum for fluid pooling in posterior fornix

Investigations :

Confirm diagnosis : Speculum / Nitrazine paper / +ve fern / Amniosure

Detect complications (infection) : CRP / TLC / DLC / ESR

US for AFI : If < 5 * oligohydramnios

Management : ANTIBIOTICS

→ IF

- < 36 wks
- NO maternal or fetal complications
- NO Ut contractions

Conservative : Antibiotics
+ Follow up maternal & fetal conditions

→ IF

- > 36 wks
- **OR** Presence of maternal or fetal complications
- **OR** Presence of Ut contractions

Antibiotics + Termination (whether VD **or** CS depending on obstetric condition)

**NO
TOCOLYSIS**

Oligohydramnios

Def: ↓ liquor < 500cc AFI < 5
or deepest pocket < 2cm

Inc : 5%

Et :

(↓ production by fetus or placenta)

- * Renal agenesis
- * Pl.insufficiency

Cl.picture:

- * Small size abdomen < amenorrhea
- * Picture of the cause (eg : PE)

Inv

US for volume / AFI / Deepest pocket

Complications:

- * Of cause (placental insufficiency)
- * Limb deformity / lung hypoplasia / amniotic band syndrome
- * Cord compression / fetal distress

Ttt:

Depends on etiology

Polyhydramnios

(Chronic hydramnios)

Def : ↑ liquor > 2liters AFI > 25
or deepest pocket > 8cm

Inc : 0.5%

Et : ↓ swallowing by fetus

or ↑ production

- * Idiopathic
- * DM (uncontrolled)
- * Oesophageal or duodenal atresia
- * Anencephaly

Cl.picture:

- * Oversize abdomen > amenorrhea
- * Pressure symptoms (as resp. embarrassment)
- * Picture of the cause (eg : DM)

Inv :

US for volume / AFI / Deepest pocket

Complications:

- * Of cause (uncontrolled DM)
- * Pressure symptoms
- * Sudden ROM , Placental abruption, Cord prolapse
- * Malpresentations , Dysfunctional labor , Atonic PPHge

Ttt:

Depends on etiology

NB: Acute hydramnios:

polyhydramnios before age of viability (24wks) associated é chromosomal anomalies , causing rapid accumulation of liquor & pressure symptoms

ttt: induction of abortion

NB : * Amniotic fluid functions : protect from infection / regulate temperature / lung expansion / limb movements / nutrition

* Amniotic fluid composition : 99% water, clear, alkaline / mainly **FETAL** contribution

Preterm labor

Definition: start of labor pains after fetal viability & < 36 wks

Incidence: 5-10%

Etiology:

- Idiopathic / miscalculation .
- Cx incompetence / septate or bicornuate uterus
- Over distended ut (polyhydramnios / MFG / fibroid ut)
- Medical / obstetric indication for termination.
- PROM / GTI / UTI / smoking / excessive physical activity
- Iatrogenic (induced PTL in complicated uncontrolled pregnancies)

Clinical picture:

→ **Symptoms :** true labor pains < 36 wks

→ **Signs:**

- General : of cause or risk factor
- Abd : true ut contractions / cause
- PV: start cx changes (dilatation / effacement)

Complications (ALL FETAL) :

→ RDS / Retinopathy of prematurity / Cerebral hge

→ Neonatal sepsis / Necrotising enterocolitis

Investigations :

→ CTG to confirm PTL

→ US : short cervical canal < 2.5 cm

→ FFN (Fetal fibronectin): from 24-32 wks

Ⓝ absent , if present (by vag swab) * 50% PTL within 2 wks

Management:

* Prophylactic against RDS: steroids 24mg IM

& delivery 24hrs after last dose :

Betamethasone (long acting) 12mg $\xrightarrow{24h}$ 12mg

Dexamethasone (short acting) 6mg $\xrightarrow{12h}$ 6mg $\xrightarrow{12h}$ 6mg $\xrightarrow{12h}$ 6mg

* IF still in latent phase (Cx < 4cm dilatation & < 50% effacement)

can use Tocolysis to delay labor till Steroids work / NICU transfer:

- 1) Ca channel Blocker (Nifedipine)
- 2) β_2 agonist (Ritodrine)
- 3) PGs synthetase inhibitor (Indomethacin) < 32wks
- 4) Anti- oxytocin (Atosiban)
- 5) MgSO₄ if < 28wks for neuroprotection
- 6) Natural Progesterone (17 OH Progesterone caproate) IM weekly

Post term pregnancy

Definition: pregnancy continue after 42 weeks

Incidence: 5-10%

Etiology:

- Miscalculation.
- Idiopathic.
- Anomalies as Anencephaly.
- Placental cause

Clinical picture:

→ **Symptoms :** of cause / may be oversized abdomen (80%)

→ **Signs :** may be normal or oversized abdomen (80%)

Complications:

→ 80% LGA (in normally functioning placenta)

→ 20% IUGR with its sequelae (in placental aging)

(postmaturity syndrome)

→ ↓ liquor / meconium stained liquor

& meconium aspiration during delivery

Investigations:

→ US : for AFI / placental grading

→ Growth curves

Management:

Assessment of fetal wellbeing (DFMC / NST / BPP / Doppler)

When to terminate:

- 42 wks
- occurrence of labor pains
- occurrence of complications

How to terminate: IOL


- Amniotomy
- Oxytocin : If Bishop > 5
- PGs : IF Bishop < 5

	0	1	2	3
Dilatation	closed	1-2 cm	3-4 cm	>5 cm
Effacement	< 30%	30-50%	50-80%	>80%
Position	posterior	mid	anterior	
Consistency	firm	mid	soft	
Station	-3	-2	-1 / 0	+1 / +2

Bishop score

Fetal growth disorders

SGA (IUGR)



LGA (Macrosomia)

Def: < 10th percentile for GA

Types:

- 1) Symmetrical 20% (fetal / intrinsic) causes
- 2) Asymmetrical 80% (maternal / extrinsic) causes

Et :

Fetal : ch.anomalies / infection in utero

Maternal : pl. insufficiency

Pathology : brain sparing effect in asymmetrical IUGR

Cl.picture :

Symptoms : small sized abdomen (< period of amenorrhea)

Signs: → General : of cause
→ Abdominal : ↓↓ FL / ↓↓ SFH (24 - 32 wks)

Inv :

- 1) To detect etiology
- 2) US for fetal growth curve assessment:

* In symmetrical IUGR → -- BPD , -- FL & --AC

* In asymmetrical IUGR → (N) BPD & --FL --AC

ttt:

according to cause

Def : > 90th percentile for GA

Et : Idiopathic / DM / Past date

Cl.picture: → **Symptoms :** of the cause / oversized abdomen

→ **Signs:** → General : of cause
→ Abdominal : ++ FL / ++ SFH

Inv :

- 1) To detect etiology
- 2) US for fetal growth curve assessment:

ttt: depends on etiology

Delivery by IOL or CS according to obstetric indication but **avoid Instrumental delivery** for fear of shoulder dystocia

IUFD

Definition: Death in utero after fetal viability

Etiology: → Idiopathic in 50%
→ Maternal as in placental insufficiency
→ Fetal anomalies / infections

Cl.picture: → ↓ or NO fetal movements
→ ↓↓ FL < period of amenorrhea


Complications : DIC if retained for weeks

ttt: → Wait for spontaneous labor pains while closely monitoring fibrinogen level , FDPs to guard against DIC
→ If anxious or start complication :
Terminate IOL OR CS according to obstetric condition

Fetal asphyxia (In utero)

Def: ↓O₂ & ↓ elimination of CO₂ * ↑ CO₂ → acidosis (PH < 7.2)

Etiology:



- 1) Maternal : uncontrolled medical disorders
- 2) Placenta : separation / insufficiency
- 3) cord: prolapsed / loops around the neck
- 4) fetus: anomalies / instrumental deliveries

Cl. Picture: (FETAL)

- 1) Abnormal CTG in assessment of fetal wellbeing:
 - Loss of beat to beat variability
 - Sinusoidal rhythm
 - Late deceleration (sp.persistent)
 - Brady < 100 OR Tachy > 160 b/min
- 2) Meconium stained liquor in cephalic presentation
- 3) Fetal scalp PH < 7.25

Management:

*** First aid measures:**

- 1) Stop oxytocin + IV fluid rehydration
- 2) Turn mother to Lt lateral position + O₂ mask
- 3) Atropine to the mother

*** If distress**

- Relieved : continue VD é continous CTG monitoring
- Not relieved : immediate delivery

If engaged presenting part
& fully dilated Cx

Forceps OR
Breech extraction

If not fully dilated

CS

Neonatal asphyxia (Post natal)

Def: ↓O₂ & ↓ elimination of CO₂ * ↑ CO₂ → acidosis in neonate

Etiology:

- persistent fetal asphyxia
- morphine given to the mother 2 – 4 hrs before delivery
- meconium aspiration
- cong anomalies of respiratory , circulatory systems
- prematurity
- birth injuries

Cl. Picture: APGAR score at
 → 1min for need of resuscitation
 → 5min for prognosis

	0	1	2
<u>Appearance</u> (color)	Blue	Trunk → pink Extremities → blue	Pink
<u>Pulse</u>	—	< 100 b/min	> 100b/min
<u>Grimace</u> (reflexes)	—	Grimace	Active cough & sneeze
<u>Activity</u> (movement)	Flaccid	Some flexion	Active movement
<u>Respiration</u>	—	Slow , irregular	Active cry

Ttt:

- Prevention → proper ANC & control of maternal diseases
 → proper intranatal care

- ttt : A (Airway & suction)
 B (Breathing & O₂ mask)
 C (Circulation CPR & warmth)
 D (Drugs) :
 - Naloxone : morphine antidote
 - NaHCO₃ : to combat acidosis
 - Adrenaline : to combat bradycardia
 - Antibiotics : in case of sepsis

* Proper use of Instrumental delivery
 * Episiotomy whenever needed
 * Proper use of morphine
 * Care during delivery of
 after coming head

Fetal birth injuries

Definition: Injuries of fetus at birth (iatrogenic)

Etiology: (Instrumental / prematurity / CPD

Types:

1) Bone injuries:

- Skull : $\pm$ ICHge*

	Subperiosteal Hge (cephalhematoma)	Caput succedaneum (scalp edema) (chignon)
Causes	Wrong application of forceps / * Depressed fracture or fissure fracture	Cervical caput: prolonged labor /cervical causes Pelvic caput : obstructed labor / CPD Artificial caput: ventouse application
When	After few hours	At birth
Shape	Overlie a certain bone Never crosses suture line Skin over it is normal	Any area of the scalp May cross the suture line Skin over it is echymotic
Fate	May be infected Calcification Hyperbilirubinemia ttt: $\rightarrow$ Expectant ttt (Antibiotics & follow up) $\rightarrow$ Measures to $\downarrow$ ICT (in cases of ICHge)	Subside spontaneously in 1-2 days

- Other bones : humerus / clavicle / spine injuries / shoulder dislocation

$\rightarrow$ ttt: Splint / slab (for long bones injury)

2) Muscle : as sternomastoid $\rightarrow$ ttt: Passive traction

3) Nerves : $\rightarrow$ Brachial plexus

- C5,6 : Erb's palsy (Policeman tip position)

- C8,T1: Klumpke's palsy (failure of grasp reflex) (Ape hand)


$\rightarrow$ Facial nerve injury : (flat nasolabial fold)

due to forceps pressure at stylomastoid foramen

ttt: Physiotherapy
(will resolve if edema)

4) Organs : liver / spleen / anus / hymen (as in breech delivery)

NB: proper management of the 2nd stage of labor will prevent most of these iatrogenic birth injuries



Erb's palsy

Klumpke's palsy

Puerperium

Definition : period of 6 wks following delivery during which all changes that occurred during pregnancy , will return back to normal.

Changes:

Uterus:

- Lochia(endometrium) : rubra → serosa → alba
- Myolysis
- Size at umbilicus on day of delivery
then SP (2wks later)
then prepregnancy size (4wks later)
- After pains: contractions that occur – specially é breast feeding – to help uterine involution
- Cx & lig back to normal (4wks later)

Vagina / Vulva :

- Reformation of rugae
- Gapping of vulva disappears

Breast :

- PRL for milk formation
- Oxytocin for milk expression
- Colostrum (++Ptns , ++ Ig , -- CHO , -- Fat)
first 3 days , then milk expressed
- lactational amenorrhea in 50% of females
- pigmentation of areola remains

vital signs :

- BP back to normal
- ++ T° , but < 38 in 1st 24hrs after labor
- Transient ++ pulse
- Diuresis (retention only with painful episiotomy)
- Constipation (due to lax muscles)
- Emotional liability
- Post partum blues are very common (needs support & reassurance) but rarely depression (needs tt) & psychosis (needs hospitalization)

PPC Program : 1week after labor , then 4-5 weeks later
to check :

- * vital signs returned back to normal
- * breast feeding & no milk engorgement
- * uterus back to normal & lochia
- * abdominal & pelvic floor excercises
- * wounds : CS or episiotomy wound
- * care for bladder & bowel by normal evacuation
- * contraceptive counselling

Puerpueral pyrexia

Definition: ++ T° ≥ 38° after 1st 24hrs of delivery , persisting for 24hrs or recurring within 24hrs till end of puerpurium.

Etiology :

- 1) Mastitis / Breast engorgement (most common)
- 2) P. sepsis (most serious)
- 3) UTI
- 4) Wound infection
- 5) Respiratory tract infections as COVID – 19 infection
- 6) DVT , Thrombophlebitis
- 7) other causes of fever as typhoid , malaria

“ Any case of puepural pyrexia should be considered p.sepsis until proved otherwise ”

Puerperal sepsis (3rd cause of maternal mortality in Egypt)

Definition : Infection of the genital tract after labor till end of puerperium

Etiology :

- **Predisposing F :**
 - General : ↓ immunity/ anemia / DM
 - Local : tears / septic conditions / instrumental delivery
 - Prolonged labor & prolonged PROM
 - Retained parts of placenta or membranes
- **Organism :** Polymicrobial (Gram -ve / +ve / anaerobes)
- **Route of infection :**
 - Ascending from vagina
 - Autogenous from elsewhere in body
 - Exogenous from attendees

Pathology :

- **1^{ry} site (symptoms immediately after delivery)**
 - * Uterus (endometritis) , Cx , Vagina , Perineal lacerations
- **2^{ry} site (late symptoms after 7-10 days)**
 - * Parametritis , salpingo oophoritis , peritonitis , pelvic thrombophlebitis (after 14 days)

Cl.picture:

- **Symptoms:**
 - * Fever/ foul smelling discharge / lower abdominal pain / ± oedematous white swollen limbs (phlegmasia alba dollens)
- **Signs:**
 - * General :Fever / tachycardia / toxic facies / dehydration
 - * Abdominal : tenderness / guarding / rebound tenderness
 - * PV: jumping sign / horse-shoe induration around Cx
 - * ± LL affection : swollen white painful limbs
 - * Septicemia in severe untreated cases

Investigations :

* To exclude other DD :

- Breast exam (for engorgement & mastitis)
- Chest X-Ray (for chest infections)
- C/S (for wound infection)
- Urine analysis (for UTI)
- Doppler US (for DVT)

* To confirm diagnosis :

- Culture & sensitivity from discharge
- Blood test (++CRP , ++ESR , ++TLC , ++DLC (shift to the left) , ++ staff / segmented ratio)
- Pelvic US

Treatment :

* Prevention :

- by proper ANC (control anemia , DM)
- 1st stage of labor :
 - Avoid prolonged labor
 - Give antibiotics in PROM
- 2nd stage:
 - Proper use of instrumental delivery
 - Proper aseptic techniques while doing episiotomy
- 3rd stage:
 - Explore placenta & memb. for any missing parts
 - Repair of any laceration under aseptic technique

* Active ttt :

- Hospitalization
- IV fluids / IV analgesics / IV antipyretics
- IV antibiotics
 - Cephalosporins for Gram +ve
 - Gentamycin for Gram -ve (Max for 3 days as they are nephrotoxic)
 - Metronidazole for anaerobes

* Special situations : (in addition to above mentioned ttt)


- Abscess : drainage
- In retained parts : Ecbolics ± D&C
- Infected wound : remove suture& drainage
- Septic thrombophlebitis : anticoagulants& immobilization
- Pelvic abscess: Fowler position ± drainage by colpotomy

Rh isoimmunization

Definition: hemolysis of fetal RBCs by maternal Ab (problem occurs when mother is Rh negative & fetus is Rh positive)

Etiology : sensitization of the mother by Rh -ve Mother received Rh+ve blood transfusion

Pathogenesis :



- 1st baby may be affected if previous maternal blood transfusion with Rh+ve blood

- Fetus may be spared if :

- 1st baby
- 2nd baby but 1st one was Rh-ve (heterozygous father)
- With ABO incompatibility
- Small amount of exposure

Effect: depends upon

- Immune system of the mother
- Associated ABO incompatibility
- She may marry Rh -ve or heterozygous Rh+ve man

Clinical picture: (FETAL ONLY)

- Hemolytic anemia (mild form)
- Neonatal jaundice : Icterus gravis neonatorum (commonest & moderate form)
± kernicterus if bilirubin > 20 mg % cross BBB
- Hydrops fetalis (severe form) → anemic HF & generalized edema (Buddha attitude)

Investigations:

- Rh blood group of mother & father
- Indirect coomb's test to mother
- Amniocentesis (if indirect coomb's > 1/16)
- US for fetal anomalies (HSM / Ascites)

Treatment:

- Prevention: Anti D (only for non sensitized Rh -ve mothers) within 72hrs of delivery of Rh+ve fetus & at 28wks GA & immediately after any procedure done during pregnancy (Abortion , Ectopic , Amniocentesis , ...)

If indirect coomb's

- ≤ 1/16 → repeat after 4 wks
- > 1/16 → amniocentesis

If ↑ bilirubin & ↓ Hb (± hemolysis)

< 34wks

Intrauterine transfusion

> 34wks

Terminate

Hb (N) 18gm% , bilirubin (N) < 2 mg / dl

Follow up with titre

After delivery: exchange transfusion for the baby with O -ve blood

Definitive ttt is plasmapheresis for maternal blood to remove antibodies (very costly, available in limited centers, doubtful prognosis)

Medical disorders during pregnancy

1) Pregnancy Induced Hypertension (PIH)

Definition : HTN + Ptnuria > 24 wks

Types :

* **Mild :** BP $\geq$ 140/90 , ptn > 300mg

* **Severe:** → uncomplicated : BP $\geq$ 160/& or 110 , ptn > 5gm (no symptoms or lab changes)
→ complicated (é symptoms or lab changes in addition to the above mentioned criteria)

Inc : 4-7% (one of the leading causes of maternal mortality)

Et : → Risk f : PG / renal disorders / SLE / APS / DM / +++ HCG (v.mole / MFG)

→ Theories & pathophysiology:

Failure of Trophoblastic invasion in spiral arterioles



Narrower vessels with ++ vascular resistance * ↓↓ **placental perfusion** → IUGR
& release of toxic mediators → Oligohydramnios
→ Ch. Fetal distress


Reaching all or any organ (liver / brain / retina / kidney / BVs)

So affecting their function

Cl.picture :

→ (S) disease of **Signs :** BP / ptn in urine / weight (due to occult edema)

→ (S)ymptoms ONLY in severe PE :



Warning signs

→ DD : of ↑ BP / ptnuria / edema

Complications : → **Maternal :** → Severe PE é symptoms or lab changes (HELLP)
Retina / liver / kidney / brain / DIC ...
→ Eclampsia (convulsions)

→ **Fetal:** IUFD

Inv : → **Maternal :** A/C ratio > 0.3 , CBC (Hb, Hct , PLT) , LFTs , KFTs ,
coagulation , fundus exam , ? CT brain

→ **Fetal :** US (growth assessment , liquor) / assessment of fetal wellbeing

Hypertensive disorders of pregnancy :

* Preeclampsia (PIH) :

HTN + Ptnuria > 24 wks GA

* Gestational HTN : HTN > 24 wks GA

* Chronic HTN : HTN < 24 wks GA

* Superimposed PE : chronic HTN + PE

* **ECLAMPSIA :** PE + Seizures

Ttt:

Mild PE :

continue to term but NOT post term
é close observation

Severe PE :

* **Complicated :** IF Symptoms or lab changes :

MgSO₄ & Immediate termination

* **Uncomplicated :** IF only BP & ptnuria
NO symptoms or lab changes:

< 36wks

> 36wks

Antihypertensive
(labetalol / methyl dopa
calcium channel blocker)
& close observation

MgSO₄
+ terminate

Till: → reach 36wks
→ labor pains
→ occurrence of complications

Eclampsia:

- MgSO₄ / Diazepam :
to control convulsions

- IV anti hypertensive as Hydralazine

- Termination of pregnancy

NB: prediction & prevention of PE :

1) Uric acid , PAPP-A & Inhibin A
for prediction

2) Uterine artery doppler

3) Prophylaxis ? LDA / Ca / Vit D

2) DM with pregnancy

Definition :

hyperglycemia / glucosuria / microangiopathy

Et:

- **GDM** : Only during pregnancy
- **Overt DM** :
 - Type I : insulin dependent
 - Type II : non insulin dependent

Risk F :

old age , multipara , H/O of IUFD , H/O of congenital anomalies

Pathophysiology :

pregnancy is **diabetogenic** due to placental anti insulin hormones (PRG , Cortisol , HPL) & insulinase enzyme

Cl.picture :

- **Maternal**:
 - ↑ infection (vulvovaginitis)
 - Abortion / PTL / PROM / Puerperal sepsis
 - Instrumental deliveries
 - ++ DKA / PE
- **Fetal** :
 - Macrosomia / polyhydramnios
 - FCA :
 - VSD (**most common**)
 - Sacral agenesis (caudal regression syndrome) (**most specific**)
 - Sudden IUFD
 - Fetal birth injuries & shoulder dystocia
- **Neonatal** :
 - RDS
 - Hypoglycemia (give IV glucose to the new born upon delivery)
 - Hypocalcemia
 - Polycythemia
 - Hyperbilirubinemia

Inv :

- **Maternal** :
 - **Screening for all pregnant ladies (24-28wks)**:
 - 50gm 1hr OGTT If ≥ 140 mg /dl
 - 100 gm 1 hr OGTT If ≥ 165 mg /dl
 - **Diagnostic** :
 - FBS > 95 mg/dl
 - 100 gm 1 hr OGTT If > 200 mg /dl
 - **HbA1c**: to assess glycemic control over past 3 months (best indicator for occurrence of fetal congenital anomalies)
- **Fetal**
 - **US**
 - 2nd trimesteric for anomaly scan
 - 3rd trimesteric for Macrosomia , Polyhydramnios
 - **Assessment of fetal wellbeing**

Treatment:

- **Diet + exercise** : if blood sugar < 200 mg/dl
- **Continue metformin** (if was given since before pregnancy)
- **Insulin** : if blood sugar > 200 mg /dl
 - $\frac{2}{3}$ dose at morning & $\frac{1}{3}$ dose at night
 - 50 : 50 (regular : NPH)
- **××× Oral hypoglycemic ××× as teratogenic & long acting so difficult to be controlled**
- **Delivery** : depends on glycemic control
 - * if good control → * wait for spontaneous labor pains
 - * if uncontrolled → * terminate after giving steroids for lung maturity.

3) UTI é pregnancy

A) Asymptomatic bacteruria

Def : > 100.000 CFU

Inc: 3-7% of pregnant females
(↑↑ in 2nd trimester due to stasis & compression)

Diagnosis : → urine analysis
→ urine culture & sensitivity

Ttt : outpatient
oral antibiotics is a must as 30 – 45 % → pyelonephritis

B) Acute Cystitis

Def : Dysuria / frequency with no fever

Ttt : as Asymptomatic bacteruria

C) Acute pyelonephritis (upper UTI)

Def : upper UTI with systemic manifestations

Et : asymptomatic bacteruria

Cl.picture: fever/ loin pain

Complications:

→ **Maternal:** pulmonary dysfunction from sepsis & anemia
→ **Fetal :** PROM , PTL , Morbidity & Mortality

Inv : → Urine analysis / Urine culture & sensitivity
→ US
→ Assessment of fetal wellbeing

Ttt: Hospitalization + IV fluids , antipyretics , analgesics
Start IV antibiotics then modify according to C/S

4) Cardiac diseases é pregnancy

Types: RHDs (developing) , CHDs (developed) countries.

Pathophysiology :

→ $\pm BP = \frac{\uparrow\uparrow CO}{\uparrow SV \times \uparrow HR} \times \downarrow\downarrow TPR$ (hyperdynamic circulation)
→ Waterhammer pulse: (++ S / D) difference
→ Apex : shifted to 4th intercostal space outside MCL
→ Split S₁ / appearance of S₃ / Systolic functional murmurs (éout thrill)
→ ECG changes due to change in heart axis

Risk F : anemia / hypertension / infection / hyperthyroidism

Clinical picture (NYHA classification) :

I : Dyspnea on > ordinary effort **II** : Dyspnea at ordinary effort
III : Dyspnea on < ordinary effort **VI** : Dyspnea at rest

Effect (complications):

→ **Maternal :** Worsen NYHA classification by 1grade
→ **Fetal :** LBW / IUGR / Fetal anemia

Inv : ECG / Echo / US for fetal biometry (SGA)

Management:

*** In pregnancy**

→ More frequent ANC
→ Guard against anemia / infection / HTN / Hyperthyroidism
→ Digitalis to be continued or started whenever needed

*** In Labor:**

→ Semisitting / O₂ mask / Analgesics / Antibiotics
→ Avoid fluid overload : care é oxytocin (has ADH like action)
→ Shorten 2nd stage by forceps / ↓ bearing down
→ Smooth VD or CS (whenever indicated)
→ In 3rd stage : * Lasix to ↓ VR & heart load * **NO Ergometrine**

*** In puerperium :**

→ Breast feeding CI in HF
→ Proper selection of contraception

5) Seizures (convulsions , Epilepsy) é pregnancy

75% remain same , 25% worsen due to metabolism of anticonvulsants.

Effect of anticonvulsants on fetus : FCA , CP/MR , - VitK (é phenytoin)

Management : → **Maternal :** * Extra folic since before pregnancy 800µg daily * Monotherapy is better é least possible dose to control seizures
→ **Fetal :** anomaly scan

6) Anemia é pregnancy

Def: ↓ Hb less than 11gm%

or less than 10.5gm% in 2nd trimester

Inc: commonest medical disorder during pregnancy

Effect of pregnancy on blood :

* Physiological (dilutional anemia)

(++ plasma > + RBCs)

* Tachycardia & Hyperdynamic circulation

* Max effect at 20 wks

Etiology :

* Nutritional (Fe deficiency anemia) **COMMONEST**

* Megaloblastic (folic A. & Vit B₁₂ deficiency)

* Hemorrhagic

(bleeding in early, late pregnancy & PPHge)

* Hemolytic (congenital or acquired)

* Hereditary (thalassemia , sickle cell anemia)

* Aplastic

Effect (complications):

→ **Maternal:** easy fatigability , PTL , PPH ,

Puerpural sepsis

→ **Fetal:** IUGR, LBW, PTL, neonatal sepsis , anemia

Inv:

→ Maternal (for iron deficiency anemia):

CBC / serum ferritin / TIBC

→ **Fetal :** Assessment of fetal wellbeing

Ttt:

→ Prophylactic

→ Mild (10-11gm/dl) : oral iron

→ Moderate (7-10gm/dl) :parentral iron

→ Severe (4-7gm/dl) / Decompensated

(< 4gm/dl): blood or packed RBCs

→ Guard against PPH / P.sepsis

→ Continue iron in puerperium

7) Thyroid disorders é pregnancy

A) physiological Goiter

Due to → ++ blood supply
→ - - iodine
→ ++ total T₃ & T₄

due to ↑ TBG

(free T₃ & T₄ unchanged)

B) Hyperthyroidism

* Graves' disease (commonest during pregnancy):
Autoimmune

* **Complications:**
Abortion , PTL , IUGR

* **Inv:** - -TSH , + + freeT₄ ,

+ +TSH receptor Ab

* **Ttt:**

Propylthiouracil is used in 1st trimester then methimazole is used for the rest of pregnancy

± βB / Steroids

× × **NO** radioactive iodine × ×

C) Hypothyroidism

Rare in pregnancy as it causes infertility & anovulation

Commonest cause:

Autoimmune

(Hashimoto thyroiditis)

Inv : + + TSH , - - freeT₄

Ttt : Eltroxine

NB:

requirements ↑↑ in pregnancy

8) Venous thromboembolism

Effect of pregnancy on blood:

→ + + clotting factors & - - fibrinolysis

→ + + platelets activation

→ Venous stasis due to pressure by gravid ut.

→ + + + thromboembolic events in pregnancy & puerperium

Risk F: > 35 yrs , multipara , obese , VV, H/O of DVT , thrombophilia , APS , CS, sepsis sp . pelvic

Cl.pict of DVT: red , hot , tender , swollen calf ms

Complications : 25% of DVT will result in PE of which 15% are fatal

Inv: → Doppler US

→ Venography (**CI** during pregnancy)

Ttt:

→ Prophylactic :

* Hydration & mobilization

* LDA

* Elastic stocking

* LMWH (clexane) single SC injection :

in high risk multifactorial , till end of puerperium

→ **Actual ttt :**

Therapeutic dose of IV heparin or LMWH & immobilization

NB: Warfarin is **CI** in 1st trimester (teratogenic)

& in 3rd trimester r (fetal ICHge)

When to stop anticoagulants before delivery?

- Aspirin 1 week before delivery

- Clexane 24hrs before delivery

9) Pulmonary Embolism

Cl.pict: Breathlessness / Hypoxia / Tachycardia / Pleuritic chest pain

Inv: → ECG changes

→ Ventilation / perfusion scan

→ Pulmonary angiography

Ttt: ICU support , O₂ Therapy ,

IV Anticoagulant (high therapeutic dose)

10) GIT disorders with pregnancy	11) Hepatic disorders with pregnancy
A) Emesis Gravidarum (NVP)	A) HBV
Reassurance / small frequent meals / ↓↓ fat & ↑↑ carbohydrate intake	DNA virus Through → Transplacental (rare) → During vaginal delivery (secretions with HBs Ag) Inv : screen for HBsAg / LFTs / PCR Ttt: → If high viral load (PCR) : ttt in 3rd trimester → VD allowed → Neonate : Ig & active immunization → Breast feeding allowed
B) Hyperemesis Gravidarum (HEG)	B) HCV
Def: Excessive vomiting in 1st trimester that affects general condition Etiology (theories) : ++ HCG / -- Vit B₁ (Thiamine) / Psychogenic Pathogenesis : → Dehydration / hemoconcentration & → Electrolyte disturbance / Starvation ketosis Complications → ↓ liver glycogen & ++ AST , ALT → Mallory Weiss Syndrome → Wernicke's encephalopathy Inv : Na⁺ / K⁺ / LFTs / Hct / chloride in urine Treatment: → Hospitalization / NPO → IV fluids & correct electrolytes → IV or PR Antiemetics as metoclopramide , meclozine , ondansetrone → Rarely termination (in severe cases of encephalopathy)	Ttt : → ttt is contraindicated during pregnancy → Avoid instrumental delivery to minimize abrasions → Breast feeding allowed except with cracked nipples
C) GERD	C) Intrahepatic cholestasis
Def : reflux aggravated by meals and recumbency Ttt: → Reassurance → H₂ receptor blocker as cimetidine Or → Proton pump inhibitor as omeprazol	Def : cholestasis & pruritis > 20 wks GA Inc : 1-4 % , Etiology : unknown ?! genetic Diagnosis : → Cl.picture : → Itching éout rash , sp. palms & soles → Jaundice (rare) → Inv: → ++ bile acids → Mild + AST, ALT , Bilirubin Ttt: → Symptomatic : cold baths , antihistaminics → Ursodeoxycholic acid tab → Termination if : reach 36 weeks / occurance of labor pains / maternal or fetal complications
	D) AFLP (Acute & may be fatal)
	Definition: fat deposition within liver cells in 3rd trimester * affection of function +++AST & ALT Incidence : extremely rare Etiology : unknown ?! error of metabolism (enzymatic) Cl.picture : → nausea , vomiting , abdominal pain , jaundice → hypoglycemia , HTN (PE) → coagulopathy , DIC Inv: ++PT , PC , INR , Bilirubin , AST , ALT , hypoglycemia ttd: prompt delivery (serious condition) & ICU admission to support general condition

Prenatal diagnosis of congenital anomalies

Definition : In utero identification of structural or chromosomal (aneuploidy / non disjunction) abnormalities in the fetus for early management (ie early induction of abortion if needed , counselling of the parents to be prepared)

Indications: High risk cases needing screening

- 1) Maternal age > 35 yrs (more risky for Down syndrome)
- 2) Early exposure to teratogens (drugs , radiation , infection)
- 3) Previous H/O of anomalies in babies or family H/O of genetic disorders

NB : If screening test is positive → * confirmatory test is needed



Screening Tests (cheap / non invasive / high sensitivity)	Diagnostic (confirmatory) Tests (accurate / invasive / high specificity)
1) Maternal serum biochemical markers : - DMT : β HCG + PAPP (11 – 13 wks) - TMT : β HCG + MSAFP + uE ₃ (14 – 16 wks) - Quadruple test : same markers as TMT + Inhibin A (16 – 20 wks) NB : <u>in Down syndrome</u> all markers ↓↓↓ EXCEPT β HCG ↑↑↑	1) Chorion Villous Sampling : (trophoblastic tissue biopsy) - vaginal (at 11wks) / abdominal (at 14 wks) - both TAS guided / abortion risk 2 %
2) US : - NT , presence of nasal bone (11-13 wks) (NT > 3mm , hypoplastic nasal bone <u>in Down</u>)	2) Amniocentesis: (Amniotic fluid & cells shedded obtained by needle US guided) - abdominal at 14-16 wks - risk of abortion 1% , infection , ROM , Pl.separation
3) Cell free Fetal DNA : (10 -11 wks) - used as screening & confirmatory - non invasive but expensive & not available in all centers - diagnostic only for 5 -12 chromosomes (including ch. 21, 13 , 18) , but not the whole 23pairs of ch. as other diagnostic tests as CVS & amniocentesis	3) Cordocentesis : (Fetal blood sample , US guided) - diagnostic & therapeutic in exchange transfusion in Rh isosensitization - abdominal at 16-20wks
	4) Advanced US (level 2) - anencephaly , cystic hygroma , skeletal anomalies (11-13 wks) - major anomalies : NTDs , skeletal , cardiac , renal & GIT anomalies , diaphragmatic & ventral hernias (18-22 wks)
	5) Cell free Fetal DNA: (as before)

NB : PGD (Pre implantation genetic diagnosis):

Done in association with IVF procedure

Single cell at 8 cell stage or dozen cells in blastocyst stage :

→ stage can be taken with no damage to fetus , to provide DNA for PCR analysis for inherited genetic disorders.

“ for diagnosis of anomalies before doing the embryo transfer & in preimplantation sex selection (in cases of sex linked anomalies)”

Prenatal diagnosis of Down syndrome	
H/O : maternal old age	
Screening	Confirmatory
10wks: Cell Free Fetal DNA 11 – 13 wks: → DMT : $\uparrow \beta$ HCG + $\downarrow$ PAPP → US: * NT > 3mm , * hypoplastic nasal bone	10wks: Cell Free Fetal DNA 11wks: Vaginal CVS 14-16 wks: Amniocentesis
14-16 wks: TMT: $\uparrow \beta$ HCG + $\downarrow$ MSAFP + $\downarrow$ uE ₃ 18-22wks: US: Other soft markers	

Fertilization , Implantation & Placenta formation

3) Fetal development

- Endoderm
- Mesoderm
- Ectoderm


é

a) Placenta formation

b) Umbilical cord


c) Extraembryonic membranes (amnion /chorion)

d) Amniotic fluid (1.5 liters , 99% water)




2) Implantation

1) Fertilization



Fetal circulation



Placenta


Definition : interface between mother & fetus

Formation : Chorion frondosum + decidua basalis

Gross: 20 cm diameter (discoid) , 500gm weight , formed of 15-20 cotyledons (unit of the placenta)


Functions:

- 1) **Transfer :**
 - **Simple diffusion :** é conc gradient as H_2O , O_2 , CO_2
 - **Facilitated diffusion :** é carrier as glucose , ketones , FA
 - **Active transport :** against conc. gradient as aa , Ca^{++} , Fe
 - **Pinocytosis :** for large molecules as LATS , Ig
- 2) **Endocrine :**
 - **HCG :** glycoptn to maintain CL of pregnancy in 1st△
 - **HPL :** ptn , main metabolic hormone , provide glucose , aa , FA to fetus
 - **Estrogen :** steroid for hyperplasia of ut ms , breast duct
↑PRL , ++ Oxytocin receptors
 - **PRG :** steroid for hypertrophy of myometrium , decidua, Breast alveoli , oedema , immunological
- 5) **Immunological :** for fetal acceptance



Abnormalities :

- 1) **In shape :**
 - **Membranacea (diffuse / large) placenta (Circumvellate)**
 - **Bilobate :** 2 lobes é tissue inbetween
 - **Bipartite :** 2 parts é membranes inbetween
 - **Succenturiate :** one large part & another small accessory lobe (may be missed & retained in delivery)
- 2) **In size & weight :** ++ in \$, hydrops fetalis , DM (> 600gm & > 5cm thickness)
- 3) **In adhesions :** accreta / increta / percreta
- 4) **In implantation :** on LUS = pl.previa
- 5) **Pl.infarcts:**
 - White : fibrin deposition / calcium deposition
 - Red : hge as in HTN
- 6) **Calcification:** with advanced aging of placenta




Umbilical cord

Definition : develops from the ventral (connecting) stalk , mesodermal in origin
measures 50cm , 1-2 cm diameter

Abnormalities :

- 1) **Insertion :**
 - Marginal (battledore)
 - Velamentous (in membranes)
- 2) **Knots :**
 - False: accumulated Wharton's jelly (no complications)
 - True :→ fetal asphyxia
- 3) **Length :**
 - Too long (> 60 cm)→ cord prolapsed & true knots , loops around neck
 - Too short (< 35 cm)→ prolonged 2nd stage of labor



4) **Vasa previa :** Vessels crossing Cx . connecting placenta to another lobe as in velamentous insertion (If bleeding→ severe fetal distress ± death)

Diagnosis of pregnancy	Physiological changes during pregnancy	ANC
1) β HCG detection : <u>* Urinary pregnancy test :</u> 1week after missed period <u>* Blood HCG :</u> Quantitative : at time of missed period $\pm$ doubling Qualitative : at time of missed period (just +ve or -ve) 2) Ultrasound : - TVS : at 5weeks (β HCG 1500 iu/ml) - TAS : at 6weeks (β HCG 6500 iu/ml) at 2.5 cm GS at 1 cm CRL GS Yolk sac or CRL +ve pulsations 5 wks 6 wks NB: If the next step is not seen : rescan after 1wk 3) Abd exam : at 2nd trimester - At 12 wks $\rightarrow$ FL at SP - At 20-24 wks $\rightarrow$ FL at umbilicus - At 36 wks $\rightarrow$ FL at XS (Leopold maneuver & FHS auscultation) When engagement occurs $\rightarrow$ FL $\downarrow\downarrow$ NB : 4) PV : is not routinely done , only in certain situation as early vag bleeding , ...	Systems: 1) Respiratory : - Hyperventilation 2) Cardiac : - Apex change - $\pm$BP = CO$\times$TPR - Bl volume $\uparrow\uparrow$ (Plasma > cells) 3) Blood: - Hypercoagulable - Stasis 4) GIT : - Morning sickness - Ptyalism (salivation) - Heart burn - Constipation 5) Urinary : - Frequency in 1st / 3rd trimesters 6) Endocrine: ++ E₂ $\rightarrow$ ++ SHBG ^ ++ Total form of hormones , but not free (active) form 2) Breast : - $\uparrow$ size / 2^{ry} areola deep pigmentation / nipple - Montgomery tubercles - colostrum 1) Genital: - Ut $\rightarrow$ UUS - LUS - Cx , vag , vulva increase vascularity - Chadwick's sign 3) Skin : - Striae gravidarum - Linea nigra - chloasma of pregnancy 4) Back : - Backache (lig relaxation) - lordosis 5) LL: - Oedema gravitational & compression	Def : program of preventive obstetric Frequency : / month in 1st 6 months / 2 weeks in 7th & 8th months / week in 9th month What to check ? Return visits $\rightarrow$ BP/ BW $\rightarrow$ Leopold maneuvers / FHS $\rightarrow$ CBC / urine analysis Booking $\rightarrow$ H/O (present /past / family) $\rightarrow$ Rh Blood group / CBC / LFTs $\rightarrow$ KFTs / FBS / urine analysis / $\rightarrow$ Hepatitis markers Supplement : 1st trimester $\rightarrow$ folic acid 2nd , 3rd $\rightarrow$ Ca, Fe , Vit Drugs & vaccines - Preferable class B medications - Only killed vaccines, toxoids and passive immunization (Ig) are given , while live attenuated vaccines are contraindicated US : - 1st T : single / living / intrauterine - 2nd T : anomaly scan - 3rd T : growth curves Special care for high risk - Elderly primi - Grand multipara $\geq$ P5 - Medical disorders - Obstetric complications

High risk pregnancies

* **Definition :** pregnancy that endangers health of mother / fetus or newborn


* **Examples :**

- 1) Elderly primi (> 35 yrs) , pregnancy > 40 yrs old , grand multipara (≥ 5 deliveries)
- 2) Maternal medical condition : uncontrolled DM , HTN , cardiac , SLE, ...
- 3) Obstetric problems: H/O of RPL, current APHge , ROM , PTL ,Placenta accreta, ...
- 4) Fetal problems : anomalies , IUGR , Multiple pregnancies , ...




* **Management :**

- More frequent ANC visits
- Delivery in specialized equipped place
- Management in pregnancy & labor accordingly

Instruments in OBGYN

	NO ANESTHESIA NEEDED				
	Uterine Sound	Cusco speculum	Sim's speculum	Volsellum	Ring forceps
			 Sim's position	 multiple toothed single toothed	
Uses	- Measures the length of the uterine cavity preliminary to IUD insertion, or any procedure as dilatation	- Used during gynecological examination to visualize the cervix and lateral vaginal walls.	- Used to retract posterior vaginal wall to expose the anterior one in case of vesicovaginal fistulae. Patient should lie in sim's position.	- Used to hold and steady the firm (gynecology) cervix for any procedure of IUD insertion.	- Used to hold and steady the soft (obstetrics) cervix in cases of early pregnancy prior to surgical evacuation as it is less traumatic than the volsellum
Comp.	Uterine Perforation	_____	_____	May be traumatic to the cervix, causing lacerations especially in single tooth volsellum	_____

Instruments in OBGYN

ANESTHESIA NEEDED				
	Dilator	Uterine Curette	Doyen Retractor	Ovum forceps
				
Uses	- Dilatation preliminary to another operation	- for endometrial biopsy (Fractional curettage) or evacuation of products of conception	- Used to retract the dissected urinary bladder down from the lower uterine segment during Cesarean section, to protect it and prevent bladder injury.	- Used to evacuate the products of conception in case of abortion - It's same as ring forceps but without a lock
Comp.	- Shock: if used without anesthesia. - Perforation of the uterus - Lacerations of the cervix - Introduction of Infection. - Anesthetic complications.	In addition to complications of dilatation, over curettage may lead to Asherman's syndrome.		



NB: In case of uterine perforation :

- Stop procedure
- Give antibiotics
- Observe vital signs
 - If normal : Discharge the patient
 - If deteriorates or intestinal contents appear through Cx : Exploratory laparotomy and proceed.

Clinical History Taking

General sheet

Personal H/O : Name / Age / Marital status / Parity / Occupation / Residency / Special habits

C/O : in patient's own words (+ duration)

Present H/O :

Gynecological sheet	Obstetric sheet	WARNING SYMPTOMS
1) Analysis of complaint (onset / course / duration) 2) Analysis of Pain / Bleeding / Mass 3) Ask about <u>etiological</u> F , symptoms & complications of your <u>DD</u> to reach a provisional diagnosis 4) Investigations & ttt done for this patient 5) Review of other systems involvement	1) LMP EDD $\pm$ GA 2) Analysis of 1st trimesteric symptoms ↳ Vomiting ↳ Bleeding ↳ Frequency of micturition 3) Analysis of 2nd trimesteric symptoms ↳ Quickening ↳ Relieve of 1st trimesteric symptoms 4) Analysis of 3rd trimesteric symptoms: ⚠ (warning symptoms) ⚠ 5) Analysis of C /O (<u>if present</u>) 6) Investigations & ttt done 7) Review of other systems	

Past H/O: Medical : medical disorders prior to C/O

Surgical : operations done prior to C/O

Family H/O : of similar condition / consanguinity

Menstrual H/O : menarche / menstrual index eg : $\frac{3}{28}$ / dysmenorrhea / intermenstrual bleeding

Obstetric H/O : F / P / A / L / GPL / history of contraceptions

F	P	A	L
> 36 wks	24-36	< 24	now

eg:

Year of pregnancy	GA at termination / delivery	Outcome	Mode of delivery / termination	place	Pregnancy complications	Postpartum complications	lactation
1999	$\pm$ 8 wks	abortion	SE	KA	NO		
2003	$\pm$ FT	L ♀	VD	home			Failure of lactation
2007	$\pm$ 28wks	SB ♂	CS	Private clinic		Wound sepsis	

IUD

OCPs

Provisional diagnosis : patient's name ثلاثي / Age / GPL / complaint in medical terms (+ duration) / provisional diagnosis / relevant medical problems / relevant surgical operations

Examples of certain sheets:

I. Bleeding sheet

Personal history : as general sheet

Complaint: Excessive or Irregular or Excessive irregular vaginal bleeding for how long

Present history:

- Analysis of the complaint:(menorrhagia, metrorrhagia, menometrorrhagia)
 - onset, course ,duration , amount of bleeding (number of pads, clots)
- Analysis of other related gynecological complaints :
 - pain (type,site,radiation,severity,.....)
 - mass (onset, course, duration,site,..)
 - discharge (amount, colour, odour ,itching)
- Analysis of the causes :
 - preceding event e.g. period of amenorrhea, I.U.D. insertion, injectables
 - contact bleeding
 - bleeding from other body orifices, ecchymosis
 - thyroid disorders
 - heart disease (dyspnea ,palpitation ,L.L.O. ,....)
 - hypertension
- Analysis of the Consequences: anemia (dyspnea, easy fatigability, blurring of vision)
- Analysis of investigations and treatment : U/S, D&C, CBC, coagulation profile
- Review of other systems.

Menstrual, Obstetric, Past, Family history, Diagnosis: as general sheet

Modifications in a case of postmenopausal bleeding

Personal History: if widow or divorced should be stated

Complaint: vaginal bleeding after years of cessation of menstruation.

Present history:

- Duration of menopause
- Symptoms suggestive of distant metastasis:
 - Lung (cough , hemoptysis,..) Liver (rt hypochondrial pain , jaundice)
 - Bone (aches and pathological fractures) Brain (projectile vomiting , headache,...)
 - G.I.T. symptoms (vomiting , constipation , bleeding per rectum)

II. Genital prolapse sheet

Personal history: as general sheet

Complaint: mass protruding from the vulva , sense of heaviness + duration

Present history:

- *Analysis of the complaint:*
 - Onset , course , duration,
 - effect of straining and lying down, (mention if the onset was following delivery)
- *Analysis of other gynecological symptoms:*
 - bleeding(menorrhagia)
 - pain (congestive dysmenorrhea)
 - discharge (leucorrhea)

N.B. these are the triad of pelvic congestion due to prolapse.
- *Analysis of the consequences :*
 - urinary symptoms: frequency, nocturia, dysuria, sense of incomplete evacuation, urine retention, loin pain, pyelonephritis, stress incontinence, inability to complete micturition except after digital reduction of the mass.
 - rectal symptoms: inability to complete defecation except after digital reduction of the mass , constipation , incontinence to flatus or stools (if associated with complete perineal tear)
 - sexual troubles: dyspareunia
 - backache (traction on uterosacral ligaments in uterine prolapsed)
- *Analysis of the Causes:*
 - precipitating factors: chronic cough , chronic constipation, obesity
 - predisposing factors: symptoms suggestive of weak mesenchyme e.g. hernia, flat foot, varicose v.
- *Analysis of investigations and treatment:* previous repair , use of pessary
- *Review of Other systems:*

In Obstetric history: It is important to ask whether her deliveries were difficult and prolonged ended with use of forceps or ventouse, delivery of macrosomic baby

Menstrual, Past, Family history, Diagnosis: the same as general sheet

III. Infertility sheet

Personal history As general sheet plus

- ask about previous marriage
- children from previous marriage
- the age of the youngest child
- husband personal history: age/ occupation/ smoking/ another marriage and the age of the youngest child from the other marriage.

Complaint: Failure of conception for years despite of regular marital life

Present history:

- *Duration of her current marriage:*
- *Analysis of the causes:*
 - **Male factor of infertility:**
 - semen analysis (results, time) - treatment (nature, duration, result)
 - previous operations (hernia, varicocele) - medical disorders and drugs.
 - **Ovarian factor:**
 - symptoms suggestive of ovulation (regular cycles ,spasmodic dysmenorrhea, premenstrual mastalgia, intermenstrual bleeding ,pain and discharge)
 - hirsutism , oligomenorrhea ,hypomenorrhea (P.C.O.)
 - symptoms suggestive of ovarian failure (hot flushes , nervousness , bony aches)
 - **Tubal factor:**
 - symptoms suggestive of salpingitis (bilateral lower abdominal pain associated with offensive vaginal discharge, fever and chills)
 - previous abdominal operations that may lead to adhesions
 - **Uterine factor :**
 - previous dilatation and curettage followed by decrease in the amount of menstrual flow (suggestive Asherman syndrome)
 - **Cervical factor:** vaginal discharge + backache, erosion, cautery, cervical amputation
 - **Sexual factor:** Frequency per week, Dyspareunia (superficial or deep), vaginismus, effluvium seminis
- *Analysis of investigations and treatment :*
 - Investigations as: hormonal profile, hystrosalpingography , sonohystrography, premenstrual endometrial biopsy, folliculometry, laparoscopy (mention the results)
 - Induction of ovulation : tablets, injections, how long,
 - History of Tuboplasty
 - Previous ART (IUI, ICSI)
- *Review of other systems:* General or Endocrine as thyroid dysfunction, DM, TB, Hyperprolactenemia

Menstrual, Obstetric, Past, Family histories and Diagnosis: as General Sheet.

IV. Primary Amenorrhea sheet

Personal history: As general sheet

Complaint: Non occurrence of menstruation Or Absence of menstruation

Present history:

- *Analysis of the complaint* a case of primary amenorrhea till age of
- *Development of secondary sexual characters :* breast development, pubic hair, axillary hair
- *Analysis of the cause:*
 - Hypothalamic cause: psychological disorders, stress, anosmia, head trauma, drugs
 - Pituitary causes: galactorrhea , symptoms suggestive of increased intra cranial tension, visual field changes
 - Ovarian causes: hirsutism, deepening of voice , pelvi-abdominal mass
 - Uterine causes: History suggestive of T.B.(night fever ,night sweat, loss of weight, loss of appetite)
 - Out flow obstruction (cryptomenorrhea): cyclic lower abdominal pain , progressive abdominal swelling, if +ve ask about urine retention .
 - General causes: thyroid dysfunction, DM, severe debilitating disease
- *Analysis of investigations and treatment:*
 - Hormonal profile, ultrasound, IVP, MRI
 - Progesterone withdrawal
 - E .+P. withdrawal
- *Review of other systems*

No menstrual or obstetric history

Family history: ask about similar condition in the family (her sisters)

Past history and Diagnosis: as general Sheet

Modifications in case of 2ry amenorrhea

Personal history: as above

Complaint: cessation of menstruation for(how long)

Present history:

- *Exclusion of pregnancy:*
 - pregnancy symptoms (nausea, vomiting, abdominal enlargement)
 - pregnancy test
- *Analysis of the last pregnancy event:*
 - post partum hemorrhage (amount , cause, blood transfusion)
 - puerperal sepsis (fever , offensive lochia)
 - in case of abortion ask about (fever , D&C, offensive discharge)
- *Hormonal treatment:* e.g. injectable contraception
- *Hypothalamic cause:* psychological troubles
- *Pituitary cause:*
 - galactorrhea
 - symptoms suggestive of pituitary adenoma (increased I.C.T. , visual field changes)
- *Ovarian cause :*
 - hirsutism , deepening of voice , pelvi-abdominal mass
 - hot flushes , nervousness, bony aches
- *Uterine cause:*
 - Symptoms suggestive of T.B.
 - history of D&C (over curettage suggestive of Asherman syndrome)
- *General cause:*
 - As 1ry amenorrhea

Menstrual history taken about menstrual condition before amenorrhea



رقم الإيداع : 2021 / 23473

I.S.B.N : 978-977-203-4062

جميع حقوق الملكية الفكرية و النشر محفوظة

هذه النسخة للتداول الإلكتروني فقط